

(12) **UK Patent Application** (19) **GB** (11) **2 370 989** (13) **A**

(43) Date of A Publication 17.07.2002

(21) Application No **0101146.9**

(22) Date of Filing **16.01.2001**

(71) Applicant(s)
BTG International Limited
(Incorporated in the United Kingdom)
10 Fleet Place, Limeburner Lane, LONDON,
EC4M 7SB, United Kingdom

(72) Inventor(s)
Amala Raman
Robert Charles Hider
Radhakrishnan Venkatasamy

(74) Agent and/or Address for Service
Richard Keith Percy
BTG International Limited, 10 Fleet Place, Limeburner
Lane, LONDON, EC4M 7SB, United Kingdom

(51) INT CL⁷
A61K 31/36 31/4025 31/45 31/4525 31/5377 , A61P
17/00 35/00


(52) UK CL (Edition T)
A5B BHA B42Y B421 B422 B423 B426 B43Y B430 B46Y
B466 B48Y B482 B52Y B522 B523 B54Y B541 B544
B55Y B550 B56Y B565 B566 B57Y B575 B58Y B586
B59Y B596 B61Y B616 B64Y B646 B66Y B664
U1S S1313 S1343

(56) Documents Cited
WO 96/25159 A1 **WO 00/02544 A2**
WPI Abstract Accession No. 1997-359094/33 &
JP 090151130 A
WPI Abstract Accession No. 1993-357134/45 &
JP 050262646 A

(58) Field of Search
UK CL (Edition S) **A5B**
INT CL⁷ **A61K 31/36 31/4025 31/45 31/4525 31/5377**
Online: **PAJ, EPODOC, WPI, CAS-ONLINE**

(54) Abstract Title
Piperine analogues for the treatment of skin conditions

(57) Compounds of formula (I) wherein m is 1,2 or 3, R¹ being an alkoxy or hydroxy group when m is 1, each R¹ being independently an alkoxy group or together being a 3',4'-methylenedioxy group when m is 2 and each R¹ being independently an alkoxy group or two R¹s being a 3',4'-methylenedioxy group and the other R¹ being an alkoxy group when m is 3, n is 0, 1 or 2, R² and R³ are both H or together represent a double bond, R⁴ and R⁵ are both H or together represent a double bond and R⁶ is a pyrrolidino, piperidino, 4-methylpiperidino, morpholino, N-monoalkylamino, N-monocycloalkylamino, 3',4'-methylenedioxybenzylamino, 2-phenethylamino or alkoxy group for use in the treatment of skin conditions. The skin conditions are either melanomas or those which are treatable by stimulation of melanocyte proliferation, such as vitiligo. The compounds are analogues of piperine and may be applied topically.



(I)

GB 2 370 989 A

Figure 1:




Figure 2:




Figure 3:




Figure 4:





Figure 5:







Figure 6.




Concentration

Figure 7.

Fig.8

Dose Response Curve



TREATMENT OF SKIN CONDITIONS

Field of the invention

5 This invention relates to the treatment of skin conditions, comprising those conditions requiring stimulation of melanocyte proliferation and to the inhibition of melanomas. The invention is of especial application to the treatment of vitiligo and skin cancer.

10 Vitiligo is a common skin pigment disorder characterised by the development of patchy de-pigmented lesions. Current treatments which include the use of photosensitisers (eg psoralens) with UVA radiation (PUVA), corticosteroids or skin grafting have low success rates and are generally accompanied by unpleasant side effects. Vitiligo has a highly detrimental impact on the emotional well-being of the sufferer, the disfiguring effects of the disease being compounded by the absence of a suitable treatment. Although vitiligo patches are not believed to contain melanocytes (pigment producing cells), a reservoir exists in hair follicles in vitiliginous skin. Thus activation of hair follicular melanocytes is a crucial process
15 in the repigmentation of vitiliginous skin.

Certain plant remedies, usually administered as mixtures of herbs or extracts, particularly those used in traditional Chinese medicine and Indian Ayurvedic medicine, have been employed for the treatment of vitiligo for a long time and in many cases have given positive results in small scale studies. Herbs such as *Psoralea corylifolia L.* and *Vernonia anthelmintica Willd.* (= *Centratherum anthelminticum Kuntze*) are well known for their use in
20 this disease. Psoralens, which are employed in the modern PUVA and khellin in KUVA therapy were originally derived from plant sources (*Psoralea corylifolia L.* and *Ammi visnaga* respectively) used in traditional remedies for vitiligo. However these therapies rely on the use of UV irradiation for their efficacy, which is associated with the aetiology of skin cancer.

25 The fruit of black pepper (*Piper nigrum L.*) and long pepper (*Piper longum L.*) are both important medicinal herbs in Ayurvedic and Unani (traditional Indian) medicine systems, in which remedies generally consist of mixtures of herbs. A wide range of the medicinal uses of black pepper have been documented by Kirtikar and Basu (Indian Medicinal Plants, 2nd Edition, Vol. 3, (1935) pages 2128 - 2135), including its use in the treatment of leucoderma.
30 Black pepper has also been implicated as a possible adjunct to *Vernonia anthelmintica* in the treatment of leucoderma (Indian Medicinal Journal, Vol. 1, 3rd Edition, (1982) 1267 - 1270).

These two herbs are employed as a constituent in many traditional herbal preparations for a variety of uses, including gastro-intestinal and skin ailments. Compositions comprising black pepper, ginger and pipali have been used in the treatment of vitiligo (Ancient Science of Life, Vol. IX, No. 4 (1990) 202 - 206); however, the specific therapeutic action of black pepper in this orally administered composition has not been established.

There is, therefore, a need for further compounds and compositions, which are able to stimulate the proliferation of melanocytes.

As reported in prior PCT Application Publication N° WO 00/02544, piperine, which is present in the fruit of *Piper nigrum*, stimulates the replication of melanocytes. The action of piperine is to increase the number of cells which confer pigmentation. Piperine is the compound (E,E)-1-[5-(1,3-benzodioxol-5-yl)-1-oxo-2,4-pentadienyl]piperidine and should not be confused with piperidine. Certain named analogues and derivatives of piperine are also mentioned in the said prior application to have such effects.

In that application, piperine is also disclosed as being able to inhibit the proliferation of melanoma cells, thus making it useful in the treatment of skin cancer.

Summary of the invention

The invention provides a method of treating a subject (human or animal) having a skin condition selected from the group consisting of (a) those requiring stimulation of melanocyte proliferation and (b) melanomas, which comprises administering to the subject, preferably to the site of the condition, an effective amount of a piperine analogue or derivative thereof, as hereinafter defined. The active ingredient may be used on its own, but is more suitably used in combination with a carrier or excipient and optionally one or more further active ingredients. Where patent law permits, the invention includes such a piperine analogue or derivative thereof for the use mentioned where such use is medical, i.e. the skin condition is a disorder.

Stimulation of melanocyte proliferation greatly facilitates the re-pigmentation of de-pigmented skin, e.g. post traumatised de-pigmented skin. The term "post traumatised de-pigmented skin" means the skin formed during the healing process that occurs after a skin trauma. De-pigmentation may arise, for example, from scar tissue formed as a result of a skin trauma such as burn or other skin lesion or may be due to vitiligo. The present invention can be used to treat any of these skin disorders in a patient.

Generally in this invention, the piperine derivative or analogue thereof may be administered by oral, topical, intravenous or subcutaneous (intra-muscular) routes but is preferably applied topically (to the area of the skin where treatment is desired).


5 The active ingredient may be formulated as a solid powder; a paste, ointment or cream; a tablet or capsules; or a solution.

The method of the invention may also be used to treat a person having a skin condition which would benefit from coloration, e.g. to enhance or promote the natural colouring of the skin. The treatment may be used for prophylactic, therapeutic or cosmetic purposes.

10 Piperine analogues or derivatives as hereinafter defined inhibit the proliferation of melanoma cells. Thus, they may also be used in the treatment of skin cancer. Another aspect of the invention therefore provides a method of treating skin cancer in a human or animal patient comprising the administration to said patient of a therapeutically effective amount of the piperine analogue or derivative thereof, as hereinafter defined.

15 The piperine analogue or derivative thereof may be administered by oral or topical routes. Suitable dosage forms may be any of those discussed above.

The formula of piperine derivatives and analogues thereof usable in this invention is given below.



(1)

20 wherein $n = 0, 1$ or 2 ;

when $n = 0$, R^2 and R^3 represent hydrogen atoms or together represent a carbon to carbon double bond;

when $n = 1$, or 2 R^2 and R^3 together and R^4 and R^5 together represent carbon to carbon double bonds, R^2 and R^3 together represent a carbon to carbon double bond and R^4 and R^5 represent

25 hydrogen atoms, or R^2, R^3, R^4 and R^5 all represent hydrogen atoms;

$m = 1, 2$ or 3 ;

when $m = 1$, R^1 represents an alkoxy group having from 1 to 3 carbon atoms or a hydroxy group;

when $m = 2$, each R^1 independently represents an alkoxy group having from 1 to 3 carbon atoms or the two R^1 's together represent a 3', 4'-methylenedioxy group;

5 when $m = 3$, two R^1 's together represent a 3'-4'-methylenedioxy group and the other R^1 represents an alkoxy group having from 1 to 3 carbon atoms or a hydroxy group;

R^6 represents a pyrrolidino, piperidino, 4-methylpiperidino or morpholino group, a N-monoalkylamino group of 4 to 6 carbon atoms, a N-monocycloalkylamino group of 4 to 7 carbon atoms, a 3'-4'-methylenedioxy-substituted benzylamino or 2-phenethylamino group or

10 R^6 represents an alkoxy group of 1 to 6 carbon atoms;

in any of its E, Z geometrically isomeric forms, but excluding the named analogues and derivatives in our prior application.

The excluded analogues and derivatives of formula (1) named in our prior application are as follows:

15 (a) n is 0, 1, or 2, m is 2, the R^1 's together represent a 3', 4'-methylenedioxy group, R^2 and R^3 , together with the carbon atoms to which they are attached form a carbon to carbon double bond and, when n is 1 or 2, R^4 and R^5 , together with the carbon atoms to which they are attached, form a carbon to carbon double bond and R^6 is piperidino, or

(b) n is 1 and (i) m is 3, the R^1 's being 3', 4'-methylenedioxy and 6'-methoxy or (ii) m is 2, the R^1 's being 3'-hydroxy-4'-methoxy; or (iii) m is 1 and the R^1 is 4'-hydroxy; and R^2 to R^6 are as defined in case (a) above, or

(c) n is 1, R^6 is piperidino, pyrrolidino, isobutylamino or methoxy and all other symbols are as defined in case (a) above, or

25 (d) n is 1, R^4 and R^5 represent hydrogen atoms and either R^2 and R^3 also do or R^2 and R^3 together with the carbon atoms to which they are attached form a carbon to carbon double bond; and m , R^1 and R^6 are as defined in case (a) above; and in all of which cases (a) to (d) the molecule is in the E,E or all E geometric configuration or in case (a) when n is 1 may be in the Z,Z, Z,E or E,Z geometric configuration.

30 Certain of the active analogues or derivatives of piperine of formula (1) are new. The present invention therefore includes such compounds, and pharmaceutical compositions containing them together with a carrier or excipient.

Brief description of the drawings

Figure 1: Plots of growth of melan-a-cells cultured in different media. Each point represents the mean and standard deviation (SD) of 6 replicates. FBS = fetal bovine serum.

5 TPA = tetradecanoyl phorbol acetate.

Figure 2: Plots of growth melan-a-cells cultured from different initial plating densities of cells. Medium was supplemented with 20 nM TPA. On day 4 the medium in the remaining plates was replaced. Each point shows mean and SD of 6 replicates.

10 Figure 3: Effect of *P. nigrum* extract on the growth of melan-a cells. Culture was maintained for 8 days. Medium and extract were replaced with fresh ones on day 4. Each point designates mean and SD of 6 replicates, except that 12 replicates were done for cells only.

15 Figure 4: Effects of *P. nigrum* extract and TPA on the proliferation of melan-a cell line. Each point shows mean and SD of 6 replicates, except that 12 replicates were done for cells only.

Figure 5: Effects of piperine and TPA on the growth of melan-a cells in the presence of RO-31-8220. n = 6 for piperine and TPA treated wells, whereas n = 12 for RO-31-8220 alone.

20 Figure 6: Effects of piperine and TPA on the growth of human melanoblasts in the presence of ET3. *P < 0.05 when compared to vehicle control (One way Anova, followed by Dunnett's t-test).

Figure 7: Effects of piperine on the growth of human melanocytes in the presence of ET1. *P < 0.05 when compared to ET1 1nM treatment (One way Anova, followed by Dunnett's t-test).

25 Figure 8: Dose response curve showing the growth of melan-a cells in presence of a compound of formula (1), RV-A01, as % of control plotted against concentration.

Description of the preferred embodiments

The following are preferred features of the compounds of formula (1) used in the invention considered alone or in any possible combination of two or more:

30 - n is 1, R² and R³ together and R⁴ and R⁵ together represent double bonds or R², R³, R⁴ and R⁵ all represent hydrogen atoms

- m is 2 or 3, two R¹'s represent 3', 4'-methylenedioxy and optionally a third R¹, representing 6'-methoxy, is also present
- R⁶ represents a piperidino, 4-methylpiperidino, pyrrolidino or morpholino group or an alkylamino group having 4, 5 or 6 atoms, preferably branched chain and especially an isobutylamino (2-methylpropylamino) group, a cycloalkylamino group having from 4 to 7 carbon atoms, especially a cyclohexylamino group, or a 3,4-methylenedioxy-substituted benzylamino or 2-phenethylamino group
- alternatively R⁶ is an alkoxy group having from 1 to 6 carbon atoms, preferably 3 to 6
- the geometric configuration at the double bonds is as in piperine (all E, E)


While the preferred meaning of R¹ is a 3',4'-methylenedioxy group, R¹ may alternatively be provided by one, two or even three groups selected from hydroxy and alkoxy having from 1 to 3 carbon atoms, preferably methoxy, e.g. as in 3'-methoxy, 4'-methoxy, 6'-methoxy and 3',4'-dimethoxy substitution of the left-hand benzene ring.

Most preferably m is 2, two R¹'s represent 3', 4'-methylenedioxy and R⁶ represents an alkylamino group having 5 or 6 carbon atoms or a cycloalkylamino having from 4 to 7 carbon atoms, especially a cyclohexylamino group, or R⁶ represents a 3,4-methylenedioxy-substituted benzylamino or 2-phenethylamino group. The symbols n, R², R³, R⁴, and R⁵ may have any of the permitted meanings, but preferably n = 1. Preferably the geometric configuration is all -E.

Specific preferred compounds of formula (1) for use in the invention are shown below. Compounds 1-14 fall within the exclusion clause above, but the substituents and variations to which they refer are illustrative: one or more such substituents or variations can be present in analogues and derivatives of piperine usable in this invention.

Variations and alterations (illustratively, all other structural features of the molecule are as in piperine unless otherwise indicated, but see above)

Variation in stereochemistry at double bonds and in extent of conjugation in chain




Compounds of formula (1) and trivial names

- 1 (E, E) - Piperine
- 2 (Z, Z) - Chavicine
- 3 (Z, E) - Isopiperine
- 4 (E, Z) - Isochavicine

- 5 3,4-dihydropiperine – Piperanine
- 6 1,2,3,4-tetrahydropiperine


Variation in separation of rings (conjugated)



Structures (all E)

- 7 $n = 0$ – Ilepcimide
- 1 $n = 1$ – Piperine
- 8 $n = 2$ – Piperettine


Alterations to nitrogen substituent




Structures (all E, E)

- $R^6 =$
- 1 piperidino - Piperine
 - 9 pyrrolidino - Trichostachine
 - 10 isobutylamino - Piperlonguminine
 - 11 methoxy - Despiperidylmethoxypiperine
 - 17 morpholino
 - 18 hexylamino
 - 19 3',4'-methylenedioxybenzylamino

Alterations to the phenyl substituent




Structures (all E, E)



- 1 3',4'-methylenedioxyphenyl; $n = 1$ - Piperine
- 12 As 1 + 6'-methoxy; $n = 1$; - Wisanine
- 13 3'-hydroxy,4'-methoxyphenyl; $n = 1$ - 4'-Methoxyisocoumaperine
- 14 4'-hydroxyphenyl; $n = 1$ - Coumaperine
- 20 4'-methoxyphenyl; $n = 0$

Alterations to connecting chain and amide group



- $R^6 =$
- 21 cyclohexylamino; $n = 1$
 - 22 cyclohexylamino; $n = 0$

The piperine analogues or derivatives can be synthesised using methods known to a skilled person (see, for example, Chapman and Hall, Combined Chemical Dictionary on CD-Rom, Release 1:1 (1997) and The Merck Index (1983), 10th edition. Publ. Merck and Co, Rahway, USA. PP. 1077-1078). Thus, they can be prepared from the appropriate acid with the appropriate connecting chain between the carboxylic acid function and the benzene ring and having the appropriate stereochemistry. Where necessary, this may be preceded or followed by reduction to reduce the double bond or bonds in the connecting chain. Methods of preparing amides and esters from these acids are illustrated by the Examples below. They may also be adapted from the references cited herein, the disclosure of which is herein incorporated by reference.

The active compounds may be formulated for topical use in the form of creams, soft paraffin or lotions. Aqueous cream BP or Yellow Soft Paraffin BP may suitably contain the active at 0.03-3.0 mg % w/w or an equivalent amount of plant extract. A suitable lotion is typically prepared from 20% glycerol and 80% ethanol in purified water and contains 0.03-3.0 mg % w/w of the active material. These topical formulations may also contain penetration enhancers such as oleic acid, propylene glycol, ethanol, urea, lauric diethanolamide or azone, dimethyl sulphoxide, decylmethyl sulphoxide, or pyrrolidone derivatives. Liposomal delivery systems may also be used.

Compositions for oral formulation include tablets or capsules containing 1.5-150 mg active or equivalent amount of plant extract.

The invention will now be described with reference to the following non-limiting examples, with reference to the accompanying tables and drawings.

Examples

25 Plant samples and preparation of extracts

Piper nigrum L. fruit (black pepper, Piperaceae), originally from India, was purchased from the Food Centre, 70 Turnpike Lane, London N8, UK. The rest of the herbs were either supplied by East-West Herbs, Kingham, Oxon, UK or by Cipla Ltd, Mumbai, India.

For the preliminary screening programme, the powdered dry herb (10 g) was heated to boiling in distilled water (100 ml) and allowed to boil for 10 min, using a hot plate as heat

source. The plant material was filtered off under vacuum through filter paper (Whatman), and the filtrate freeze-dried.

Cell culture experiments

5 Microplate culture and sulforhodamine B (SRB) assay

Cells of mouse melan-a cell line (passage number 18-24), a first known line of non-tumorigenic pigmented mouse melanocytes were maintained in a flask (Costar, Cambridge, MA, USA) using RPMI 1640 (ICN, Costa, Mesa, CA, USA) as a basic medium. For microplate proliferation assays, subconfluent melan-a cultures were trypsinized (0.25% trypsin at 37°C for 5-10 min) and inoculated with a repeater-pipettor (Finn pipette, Labsystems, Finland) into 96-well microtiter plates (Costar, Cambridge, MA, USA). The plates were incubated at 37°C in a 10% CO₂, 90% air humidified atmosphere for the stated length of time. At the end of the incubation, an SRB assay was performed. Briefly, cells attached to the bottom of the plate were fixed by addition of cold trichloroacetic acid (TCA, 4°C, Aldrich, Dorset, UK) on the top of the growth medium (final TCA 20% w/v). The plate was placed at 4°C for 1 hour before being gently washed five times with tap water. It was allowed to dry in air, or aided with a hair dryer to speed up the drying process, then 50 µl of 4% w/v SRB dissolved in 1% acetic acid in water was added to each well for 30 min. At the end of the staining period, unbound SRB was removed by washing 4 times with 1% acetic acid. The plate was air dried again, and 150 µl of 10 mM aqueous Tris base (Sigma-Aldrich Co. Ltd, Irvine, UK) was added into each well to solubilize the cell-bound dye. The plate was shaken for 15 min on a gyratory shaker followed by reading the optical density (OD) at 550 nm in a microplate spectrophotometer (Anthos Labtec HT3, version 1.06)

25 Example 1

Optimisation of incubation conditions - FBS concentration and cell seeding density

Prior to testing the herbal extracts, optimal culture conditions were established. The variable factors regarding incubation conditions include foetal bovine serum (FBS) concentration, initial cell seeding density and incubation period. To determine optimum FBS concentration, 1, 2, and 5% FBS were used to culture the melan-a cell line, the growth pattern with each concentration of FBS was monitored by SRB assay. For the determination of

optimum cell seeding density, a series of initial seeding density of 0.15 to 1.2×10^4 cell per well of melan-a cells were plated into 96-well plates with 5% FBS and 20 nM tetradecanoyl phorbol acetate (TPA) supplemented growth medium. The growth pattern was monitored with SRB assay at daily intervals. The culture was extended to 8 days; on day 4, the medium in the remaining plates was replaced.

Results

The effect of FBS concentrations on melan-a growth

The optimal condition for the negative experimental control, is that cells neither grow too fast nor decline dramatically. Rapid growth might mask any subtle stimulatory effect brought about by the herbal extracts, whereas a dramatic decline in cell numbers indicates unfavourable culture conditions for cell survival, which could lead to cell damage. Figure 1 shows the growth curves of melan-a cell line at three different concentrations of FBS. Neither 1% nor 2% FBS supplemented medium was able to maintain cell survival; cell numbers declined significantly in 4 days of culture. However, 5% FBS was capable of keeping melan-a cell line alive with only a small increase in cell numbers observed over 4 days. TPA (20 nM) was able to cause further proliferation in the presence of 5% FBS indicating that cells were capable of responding to mitogenic stimuli at 5% FBS. Morphological observations under a microscope revealed that with 1% and 2% FBS supplemented medium, cell bodies were round, lightly pigmented with few dendritic processes and the culture displayed an ageing growth pattern. However in 5% FBS, cells possessed more melanosomes and some short dendrites without an ageing appearance. Therefore 5% FBS was used throughout in the herbal screening experiments.

Growth curves of melan-a cell line with various seeding densities

In Figure 2, growth curves over 8 days with different initial cell numbers were plotted to elucidate the melan-a cell line's growth pattern in 96-well plates in the presence of 5% FBS and 20 nM TPA. The optimal initial plating density together with proper harvesting time was determined. All of the initial plating number of cells showed a net growth in the presence of TPA and 5% FBS supplemented medium, although the higher plating density of 1.2×10^4 cells/well depleted the growth medium on day 3 of culture and the cells ceased to grow until

the medium was replaced. With the lower plating densities (2 - 4 x 10³ cells/well) the SRB assay OD readings remained relatively low after 8 days' culture. The initial plating density of 6 x 10³ cells/well exhibited exponential growth, and after 4 days of culture, the OD reading increased to a value of about 0.4. Since the higher OD values are associated with greater precision and accuracy, it was determined that the initial inoculation of 6 x 10³ cells/well was the optimum density for the herbal test experiment. For the simplicity of the experiment, harvesting time was day 4 since the cells at this stage was not confluent and after 4 days, growth medium tended to become depleted and replacement was necessary for the further growth.

10

Example 2

Preliminary herbal screening experiments

Melan-a cells were seeded at a density of 6 x 10³/100µl/well in standard medium supplemented with 0 nM TPA and 5% FBS. After 4 hours of incubation, herbal extracts, which were reconstituted in growth medium and sterilised by filtration (pore size 0.2 µm), of different concentrations was added into each well. Final concentrations of plant extract were 0 (negative control), 10, 100 and 1000 µg dry extract per ml. 6 replicate wells were used for each concentration tested. The negative control (12 wells), positive control (20 nM TPA, 6 wells), and test wells were all in the same 96-well plate. The culture was terminated after 4 days and SRB assay performed according to the methods given above.

20

Results

The effect of 30 herbal extracts on the proliferation of melan-a cell line

Table 1 shows the results of the preliminary screening of 30 aqueous herbal extracts on the proliferation of melan-a cell line. Crude extracts of *Astragalus membranaceus* (Fisch.) Bunge, unripe *Citrus reticulata* Blanco, *Dictamnus dasycarpus* Turcz., *Ophiopogon japonicus* (Thunb.) Kergawe, *Piper nigrum* L., *Poria cocos* (Schw.) Wolf and *Tribulus terrestris* L. were observed to stimulate melanocyte proliferation, sometimes even at the lowest dose level of 10 µg/ml. Other extracts either had no significant effect or were cytotoxic. Among these positive responses, that of *Piper nigrum* L. extract at 0.01 and 0.1 mg/ml was the most pronounced. *Piper nigrum* extract at these two concentrations not only strikingly enhanced cell growth, but

30

this extract also altered the cell morphology. In the presence of *Piper nigrum* extract, the cellular bodies were smaller, with more and longer bipolar or polydendritic processes, an effect similar to that observed with TPA.

5 Example 3

Repeats of the tests on *Piper nigrum* extract on the melan-a cells

A newly prepared *Piper nigrum* fruit extract was tested on a new batch of melan-a cell line with the culture in microplates extended to 8 days. The effects of *Piper nigrum* extract on the growth of melan-a cell line were evaluated by SRB assay.

10

Results

Repeats of the tests of *Piper nigrum* extract on melan-a cells

In the light of the positive results from the preliminary experiment, further investigations on *Piper nigrum* extract were carried out. Figure 3 shows that the result of the significant proliferant effect brought about by the *Piper nigrum* extract was even more marked on the extension of the incubation period to 8 days of culture, the growth was 272% of the control (cells only). Microscopically, the morphology of the cells was altered as those seen in the preliminary experiments.

15

20 Example 4

Confirmation of the proliferant effect of *Piper nigrum* by haemocytometer counting

Melan-a cells were plated in petri dishes (Ø35 mm, Nunclon, Denmark) with a plating density of 2×10^4 /ml and *Piper nigrum* extract at concentrations of 0.01 and 0.1 mg/ml. A negative control (cells in medium only) and positive TPA (20 nM) control were also set up.

25 After 4 days the cells in each dish were harvested and counted with haemocytometer.

Results

Confirmation of the proliferant effect of *Piper nigrum* by haemocytometer counting

SRB assay indirectly estimates cell number through protein staining and spectrophotometric measurement. To confirm if *Piper nigrum* extract stimulates melan-a cell proliferation, a direct cell counting with haemocytometer method was employed. Table 2

30

shows the cell numbers in the presence of *Piper nigrum* extract and 20 nM TPA. Cell number under the influence of *Piper nigrum* extract at 0.01 and 0.1 mg/ml were increased significantly compared to control, but less than that with 20 nM TPA. This result is consistent with the finding in 96-well microplate SRB assay.

5

Example 5

Effect of Piperine on the growth of melan-a cell line

Piperine (Sigma-Aldrich Co. Ltd, Irvine, UK) was dissolved in MeOH, sterilised by filtration through a membrane (pore size 0.2 µm) and diluted with standard growth medium.

10 The final concentrations in culture were 0.1 and 1 µM. A separate experiment (data not shown) showed that the concentration of MeOH present in these experiments was not toxic or proliferant to the cells.

Results

15 The effect of piperine on the proliferation of melan-a cell line

The effect of this compound on melan-a cell line is shown in Figure 4. Piperine at the two concentrations tested significantly stimulated melan-a proliferation. This compound brought about morphologic changes to melan-a cells, with smaller cell bodies, more and longer cellular dendrites, resembling those alterations induced by *Piper nigrum* extract and
20 TPA. This indicates that piperine is an active principle responsible for the observed proliferant effect of *Piper nigrum*.

Example 6

Test of piperine on different cell types to determine its specificity

25 In order to determine the specificity of piperine, a panel of different cell types were employed to facilitate this investigation. These included melan-a, melan-c, SVK14, CSM, XB2, SC1, B16F10, IM9, CACO2, Swiss 3T3 cell lines and normal human lymphocytes. TPA
- (20 nM) was also tested on these cells. Table 3 shows the biological origin of the cells and an outline of the cell culture protocols.

30

Results

The effects of piperine and TPA on the growth of a panel of cell types.

From Table 4, it can be seen that piperine has a highly selective effect on the growth of a panel of cell types, since it only stimulates the mouse melanocytes (melan-a, melan-c), human melanoblasts (FM21E), human foetal melanocytes (FM 21E) and the mouse fibroblast SC1 cell lines at the concentration tested. The SC1 cell line may have a particular sensitivity to TPA due to the way in which it has been derived, i.e. it has been cultured in the presence of TPA. However, piperine has either no effect or a cytotoxic effect on other cells. This result implies that piperine may have desirable specificity index for the proliferation of melanocytes in culture and is not a general mitogen. In our experimental system, TPA, a well known PKC activator and a tumour promoting agent, had similar effects to piperine on all cell types tested, except that TPA strikingly stimulated human lymphocyte and 3T3 fibroblast proliferation whereas piperine obviously lacked such an activity. Piperine seems to be a less potent stimulant than TPA.

15

Example 7

Mode of Action: effect of RO-31-8220 on the growth of melan-a cells with piperine and TPA

Melan-a cell line cultured with piperine 1 μ M and TPA 20 nM separately was set up in a 96-well plate, 1 μ l of different concentrations of RO-31-8220 (Calbiochem-Novabiochem) in DMSO was introduced with a micro-syringe into the wells to make up the final RO-31-8220 concentrations of 0 (control), 0.1, 1, 5, 10, 100 nM, with final DMSO concentrations smaller than 0.01% v/v, at which the DMSO showed neither toxic nor proliferant effect to the cells in a separate experiment (data not shown). 6 replicate wells were used for each concentration. The culture was incubated for 4 days before it was terminated and processed with SRB assay to evaluate the growth of melan-a cells.

25

Results

Mode of Action: effect of RO-31-8220 on the growth of melan-a cells with piperine and TPA

Figure 5 shows the effect of RO-31-8220 on the survival and growth of melan-a cell line in the presence or absence of piperine and TPA. RO-31-8220 alone did not have significant cytotoxic effect to the cells at the concentrations up to 100 nM. However, the

30

proliferant effects of piperine, and TPA (as indicated by the Y axis values) on melan-a cells were effectively inhibited by the presence of RO-31-8220 at the concentrations of 0.1 - 100 nM. It thus appears that piperine and TPA exert their proliferant effects through the activation of PKC cell signalling pathway.

5 The selectivity of piperine on the growth of a panel of cell types has also been tested. It was found that piperine possessed a fairly high specificity and selectivity towards melanocytes, since it significantly stimulated the growth of melan-a, melan-c and FM21E melanoblasts and FM21E melanocytes in culture, whereas it did not stimulate all other cells apart from a TPA-sensitive fibroblast cell line. Piperine was observed to have inhibitory
10 effects on B16 mouse melanoma cell line which is syngeneic with melan-a cells. Thus piperine may be a specific stimulant for the proliferation of melanocytes in vitiliginous skin without the risk of stimulating melanoma cells.

Example 8

15 Experiments on human melanoblasts in culture

Human melanoblasts in culture in this experiment were established from human foetal skin. Subconfluent melanoblasts maintained in MCDB 153 medium supplemented with 10% FBS, 10 ng/ml stem cell factor (SCF) and 1 nM endothelin 3 were subcultured and inoculated
20 into 96-well microplate with 6×10^3 cells/100 μ l/well. After incubation in the 10% CO₂, humidified atmosphere, at 37^oC for 3 - 4 hours to allow the attachment of the cells on the plate, piperine dissolved in MeOH and water was added into the wells. The final concentrations of piperine were 1, 5, 10, 100 μ M, with TPA (20 nM) as positive control. Six replicates were used in each group of treatment, with 12 wells used for vehicle control. The incubation was conducted for 5 days before cells were harvested by fixing with cold
25 trichloroacetic acid (TCA, at 4^oC, final concentration 20% v/v), and evaluated for cell number using an SRB assay. One way ANOVA and Dunnett's t-test was employed to test the significance of any differences between treatment groups and vehicle control. Growth in the presence of piperine and TPA was expressed as % of control incubations containing no piperine or TPA. The experiments were repeated using melanoblasts from 3 different donors.

30

Results

Figure 6 shows the effect of piperine on the growth of human melanoblasts *in vitro*. Piperine at the concentrations of 1, 10, 100 μM was found to cause significant stimulation to human melanoblasts in a dose response manner, with 34% more cell yield compared to vehicle control when the culture was exposed to 100 μM piperine in culture for 5 days. TPA, a well-known melanocytic growth-stimulating agent, was also able to cause significant cell growth at tested concentrations, with over 50% of more cell yield observed when the culture was exposed to 20 nM for 5 days. In the other repeated experiments, piperine was consistently observed to induce significant cell growth at the concentrations ranging from 5 - 100 μM ; these stimulatory effects were generally less than that of TPA. Morphologically, in the presence of piperine, melanoblasts appeared to be more dendritic and the cell bodies were flatter and smaller.

Example 9

15 Experiments on human melanocytes in culture

Human melanocytes used in this experiment were derived from induced differentiation of human foetal melanoblasts. The key character of human melanocytes that is different from its precursor melanoblasts is their ability to synthesise melanin. Melanin is a valid marker for melanocytes. The cell pellet of human melanocytes exhibits a characteristic brown to black colour, whereas human melanoblasts cannot produce melanin thus devoid of brown or black colour in the cell pellet.

Two protocols were employed for the experiments on human melanocytes in culture. The first employed 24-well plates and evaluated cell number with SRB assay. The second employed petri dishes and cell number was counted with a haemocytometer chamber.

For the first protocol, subconfluent human melanocytes maintained in a $\text{Ø}100$ mm petri dish were subcultured into two 24-well plates (Falcon) using basic culture medium of RPMI 1640 supplemented with FBS (10%), bFGF (100 pM), CT (1 nM) and endothelin 1 (1 nM). The initial plating density was 20,000 cells/cm² (38,200 cells/well) with each well containing 1000 μl medium. After incubation in a 10% CO₂, humidified atmosphere, at 37°C for 2 - 3 hours to allow the attachment of the cells, piperine in 500 μl medium was added into wells to made up final concentrations of 0, 1, 5, 10 and 100 μM . Cells only in the medium

with above supplement lacking of endothelin 1 were also set up as negative control. Six replicates were used in each group of treatment, and culture was incubated for 5 days before the cells were harvested by fixing with cold TCA (final concentration 20%) and processed with SRB assay. The solubilized SRB dye solution was transferred to a 96-well plate for optical density reading.

For the second protocol, subconfluent human melanocytes were subcultured in a Ø60 mm petri dishes (28 cm², Falcon) with RPMI 1640 basic medium supplemented with FBS (10%), CT (1 nM), bFGF (100 pM) and endothelin 1 (1 nM). The initial plating density was 10,000 cells/cm², with 5 ml medium per dish. Cells were incubated for 2 - 3 hours in 10% CO₂, humidified atmosphere, at 37⁰C, followed by addition of piperine solution in to the dishes, making the final concentrations of 0, 1, 5, 10 and 100 µM. Cells in the above supplemented medium lacking endothelin 1 were also set up as a negative control. Three dishes were used for each treatment and the culture was maintained for 5 days before cells were harvested with trypsinisation and counted with a haemocytometer chamber. For melanin production experiment, the harvested cells were centrifuged and pelleted. After carefully removing the medium, NaOH (1 M) was used to solubilized the cell pellets and optical density read at 475 nm in a Perkin-Elmer UV spectrophotometer (model UV/VIS Lambda 2). The melanin content was calculated by using a regression equation $y = 0.005 + 0.005x$ corresponding to the calibration curve for synthetic melanin.

20

Results

Figure 7 delineates the effects of piperine on the growth of human melanocytes cultured in 24-well plate. Piperine at the concentrations of 5 and 10 µM markedly stimulates the growth of these pigmented cells, with 36% more cells yielded when the culture was under the influence of 10 µM piperine for 5 days. However, at 100 µM, piperine exerted inhibitory effect on the growth of these cells. In addition, in the presence of 1 nM endothelin 1, TPA at 20 nM was not able to stimulate cell growth in our culture system, a result that is of great difference with that observed in human melanoblasts.

Table 5 shows the effects of piperine on the growth of human melanocytes cultured in petri dishes. It is conspicuous that in the presence of ET1 (1nM), piperine at the concentrations of 5 and 10 µM significantly stimulated the growth of human melanocytes,

30

with cell number over twice as many as that of ET1 (1nM) control when this melanocyte cell type was exposed to 5 μ M piperine for 5 days. This result was consistent with that obtained from the 24-well plate experiments, and it served to confirm that the stimulatory effects observed by SRB assay were indeed due to increased cell number rather than augmentation of protein production alone.

5

TABLES

Table 1. Preliminary screening of 30 herbal aqueous extracts on the proliferation of melan-a cell line detected with SRB assay after 4 days culture.

Names of herbs	Plant part	Cell number (% of control) after 4 days incubation when grown in the presence of extract at:		
		1 mg ml ⁻¹	0.1 mg ml ⁻¹	0.01 mg ml ⁻¹
plants with a significant stimulatory effect				
<i>Astragalus membranaceus</i> (Fisch.) Bunge	Root	163.2*	123.6*	105.6
<i>Citrus reticulata</i> Blanco (<i>Qing Pi - unripe</i>)	Peel	16.0	138.5*	127.6*
<i>Dictamnus dasycarpus</i> Turcz.	root bark	105.0	159.4*	98.0
<i>Ophiopogon japonicus</i> (Thunb.) Kergawe	Root	127.8*	126.5*	108.4
<i>Piper nigrum</i> L.	Fruit	11.5	215.4*	151.3*
<i>Poria cocos</i> (Schw.) Wolf (fungus)	Sclerotium	79.0	134.6*	128.8*
<i>Tribulus terrestris</i> L.	Fruit	80.7	136.1*	142.2*
plants with no significant stimulatory effect				
<i>Angelica dahurica</i> (Fisch.) Benth. & Hook.	Root	50.4	118.3	107.0
<i>Chaenomeles lagenaria</i> (Loisel.) Koldz.	fruit	57.1	74.5	99.0
<i>Citrus reticulata</i> Blanco (<i>Chen Pi - ripe</i>)	Peel	34.6	101.1	81.1
<i>Corydalis bulbosa</i> D.C.	Root	91.8	101.2	92.9
<i>Curcuma longa</i> L.	Root	84.1	104.8	108.3

<i>Cyperus rotundus</i> L.	Rhizome	27.5	52.8	55.8
<i>Cornus officinalis</i> Sieb. et Zucc.	Fruit	30.4	92.1	101.6
<i>Gentiana scabra</i> Bunge	Root	42.2	107.4	108.6
<i>Ligustrum lucidum</i> Ait.	Fruit	97.6	58.1	98.4
<i>Lithospermum erythrorhizon</i> Sieb. et Zucc.	Root	43.8	103.8	111.3
<i>Notopterygium incisum</i> Ting	root / rhizome	18.1	97.4	94.8
<i>Paeonia lactiflora</i> Pall.	Root	31.8	62.2	100.7
<i>Paeonia suffruticosa</i> Andr.	Root	53.2	72.0	132.8
<i>Picrorhiza kurroa</i> Royle ex. Benth	Rhizome	42.5	77.5	90.0
<i>Platycodon grandiflorum</i> (Jacq.) A. DC.	Root	35.1	94.1	96.8
<i>Plumbago zeylanica</i> L.	Root	30.2	103.9	114.1
<i>Polygala tenuifolia</i> Willd.	Root	12.7	43.7	79.6
<i>Ramulus mari</i> (insect)	Whole	41.1	87.1	89.4
<i>Siesgesbeckia pubescens</i> Makino	Herb	17.0	40.8	51.7
<i>Spirodela polyrrhiza</i> (L.) Scheid	Herb	100	79.3	96.6
<i>Trichosanthes kirilowii</i> Maxim	Root	112.9	108.1	116.1
<i>Tripterygium wilfordii</i> Hook.	Root	89.8	36.7	63.3
<i>Zingiber officinale</i> Roscoe	Rhizome	7.9	105.8	90.6

*P < 0.01 compared with vehicle treatment (one-way ANOVA, followed by Dunnett's t-test).

Table 2. Effects of *Piper nigrum* extract on the proliferation of melan-a cells counted with haemocytometer

Treatment to cells	cell number (x 10 ⁻⁴ /ml)
Control	2.02
20 nM TPA	5.0*
<i>Piper nigrum</i> at 0.01 mg/ml	3.06*
<i>Piper nigrum</i> at 0.1 mg/ml	3.13*

*P < 0.01 compared with vehicle treatment (one-way ANOVA, followed by Dunnett's t-test).

Table 3. Biological origin and the culture conditions of a panel of different cell types used in selectivity experiment.

Cell name	biological origin	optimum culture conditions		
		FBS	Medium	incubation (4days)
Melan-a	normal epidermal melanoblasts from embryos of inbred C57BL mice	5%	RPMI1640	37°C, 10%CO ₂
Melan-c	albino embryos of outbred LAC-MF strain mice	10%	RPMI1640	37°C, 10%CO ₂
FM21E melanoblast	human foetal melanoblasts from epidermis (strain 21)	10%	MCDB153	37°C, 10%CO ₂
FM21E melanocyte	Human melanocytes derived from FM21E melanoblasts		RPMI1640	37°C, 10%CO ₂
SVK14	human keratinocytes	10%	DMEM	37°C, 10%CO ₂
CSM14.1.4	neuronal cells from mesencephalin of rat	10%	DMEM	34°C, 5%CO ₂
SC1	Fibroblastoids from neonatal murine skin	10%	DMEM	37°C, 10%CO ₂
XB2	murine keratinocytes	10%	DMEM	37°C, 10%CO ₂

B6F10	mouse melanoma	5%	RPMI1640	37°C, 10%CO ₂
CACO2	human colon cancer	10%	RPMI1640	37°C, 10%CO ₂
IM9	human lymphoblastoid B cells	10%	RPMI1640	37°C, 5%CO ₂
Swiss 3T3	mouse fibroblasts	10%	DMDM	37°C, 5%CO ₂
Human lymphocytes	healthy human blood samples	10%	DMEM	37°C, 5%CO ₂

Table 4 Effects of piperine and TPA on the growth of a panel of cell types. (see Table 3 for details of cells)

Cell type	cell number as a % of control									
	piperine at the concentration of (μM)					TPA at the concentration of				
	0.01	0.1	1	10	100	20nM	200nM	200nM	200nM	200nM
Melan-a	ND	130*	169*	153*	ND	295*	ND	ND	ND	ND
Melan-c	109	208*	198*	119*	137*	186*	222*	222*	222*	222*
FM21E Melanoblast	ND	101	101	119*	134*	153*	ND	ND	ND	ND
FM21E human melanocytes	ND	ND	98	143*	75*	98	102	102	102	102
SVK14	97	101	92	84*	23*	71*	66*	66*	66*	66*
CSM14.1.4	93	94	95	89	64*	85*	76*	76*	76*	76*
SC1	191*	178*	175*	204*	190*	178*	199*	199*	199*	199*
XB2	80	90	86	90	42*	96	99	99	99	99
B16F10	71*	64*	47*	33*	0*	35*	55*	55*	55*	55*
CACO2	103	99	102	81*	34*	95	90	90	90	90
IM9	ND	101	103	69*	ND	ND	ND	ND	ND	ND
Swiss 3T3	113	104	106	102	51*	185*	207*	207*	207*	207*
Human Lymphocytes	ND	93	93	ND	ND	282*	ND	ND	ND	ND

* $P < 0.05$ compared to treatment with vehicle alone (one-way ANOVA followed by Dunnett's t-test); ND = not done.

Relative standard deviations of all values were less than 10% of the mean

Table 5: Effects of piperine on the proliferation and melanin production of human melanocytes cultured in petri dishes

Treatments	Cell no.(x 10 ⁴) ± SD after cultured for 5 days	% of control (ET1 1nM)	OD reading ± SD at 475 nm	Melanin content/10 ⁴ cells
Cells only	17.71 ± 6.16*	49.1	0.026 ± 0.0049*	0.23 ± 0.001 µg
ET1 (1Nm)	36.04 ± 6.16	100.0	0.087 ± 0.044	0.46 ± 0.01 µg
ET1 (1nM) + piperine (1 µM)	60.83 ± 16.78	168.8	0.134 ± 0.014*	0.42 ± 0.03 µg
ET1 (1nM) + piperine (5 µM)	78.96 ± 5.63*	219.1	0.137 ± 0.0085*	0.334 ± 0.01 µg
ET1 (1nM) + Piperine (10 µM)	64.79 ± 13.47*	179.8	0.139 ± 0.028*	0.41 ± 0.07 µg
ET1 (1nM) + piperine (100 µM)	61.04 ± 10.04	169.4	0.144 ± 0.0046*	0.46 ± 0.001 µg

*, * P < 0.05 when compared to ET1 (1 nM) control (one way ANOVA, followed by Dunnett's t-test).

Example 10

Derivatives of Piperine

1.0 Introduction

5 Vitiligo is defined as a circumscribed, acquired, idiopathic, progressive hypomelanotic skin disorder which is characterised by the development of patchy depigmented macules due to progressive loss of melanocytes which is often familial with lack of established aetiology.

10 Various piperine derivatives of formula (1) were synthesised and tested for melanocyte (mouse melan-a) proliferant activity *in-vitro*. Cells were incubated with the test compound for 4 days, as previously for mouse cells but with 10% FBS, after which the sulphorhodamine-B (SRB) assay was performed to determine cell number SRB uptake was measured as optical density at 550nm. The control assay was carried out on cells incubated without test compound. There were 2 or 3 series of experiments, each of which consisted of six replicate experiments. The results are tabulated below.

15

1.1 Percentage cell growth (A)

Percentage cell growth was obtained with a given compound calculated as **(optical density in the presence of the compound/ control optical density) x 100.**

1.2 Relative activity to piperine

20 Melan-a cell proliferant activity for tested compounds was compared with that obtained with piperine. Percentage stimulant activity is (A-100) where A stands for piperine or a test compound's percentage cell growth (see 1.1). All figures are given with Standard Error of the Mean.

25 Relative activity to piperine was calculated as **(A-100) compound / (A-100) piperine).**

Interpretation of the relative active value is as follows

- 30
- < 0 - Inhibition of cell growth
 - 0 - No effect (equal to control)
 - 0-1 - Stimulant but weaker effect than piperine
 - 1 - Equal stimulant effect to piperine

> 1 - Stimulant and stronger effect than piperine

1.3 Dendricity

5 Effect on dendricity of melan-a cells by the test compounds was by observation under microscope. Dendricity is relevant to vitiligo since normal skin melanocytes have dendrites, but in vitiligo the melanocytes seem to lose these before they disappear from the patches.

1.4 Synthesis of piperine analogues

10 Analogues of piperine were synthesised using methods described in the literature, adapted from the literature or devised in the inventors' laboratory. Structures of compounds were verified using NMR, MS, IR spectroscopy and melting point. Unless a synthetic method is given, reagents and reactants were purchased from Sigma Aldrich.

1.5 Results

15 Table 6 presents an overall summary of the results appearing in detail in other Tables which follow. Tables 7-12 relate to results at a single concentration of test compound (10 μ M). They are followed by data showing results at other concentrations. Many compounds showed a "cross-over" effect in which the test compound was less
20 active than piperine at 10 μ M but more active at 50 μ M. This is illustrated for one compound (RV-A01) in Figure 8 of the Drawings.

Table 6: Overall Summary of Results





Test CPD.	Change with Respect to Piperine	Active at 10 μ M?	Active at higher conc. (μ M)?	More active than piperine at higher conc. (μ M)?
Vary amide (amino group listed below)				
RV-A01	Pyrrolidino	Yes	25, 50	50
RV-A02	Morpholino	Yes	25, 50	50
RV-A04	3,4-methylenedioxy-benzylamino	Yes ^a	50	50
RV-A05	Hexylamino	Yes	25, 50	No
RV-A06	Isobutylamino	Yes	25	No
RV-A07	Methylamino	Yes	No	No
RV-A08	Ethylamino	Yes	No	No
RV-A09	Isopropylamino	Yes	No	No
RV-A10	Cyclohexylamino	Yes	50	50
RV-A11	Butylamino	Yes	50	50
Shorten connecting chain by 2 C-atoms (1 double bond)				
RV-B01	Piperidino	Yes	25, 50	50
Shorten connecting chain by 2 C-atoms (1 double bond) and vary amino part of amide group				
RV-B02	Pyrrolidino	No ^b	Not done ^d	Not done
RV-B03	Morpholino	No	Not done ^d	Not done
Replace amide by ester (alkyl group listed below)				
RV-AB1	Methyl	Yes	50, 100	50, 100
RV-AB2	Ethyl	No ^c	Not done	Not done
RV-AB4	Isopropyl	Yes	50	50
RV-AB5	Propyl	Yes	50	50, 100 ^e
RV-AB6	Butyl	Yes	50, 100	50, 100
Shorten connecting chain as above and replace amide by ester (alkyl group listed below)				
RV-BB1	Methyl	No ^b	Not done	Not done
Reduce double bonds in connecting chain, making it saturated				
RV-C02		Yes	25, 50	No
Reduce double bonds in connecting chain and shorten it by 2 C-atoms				
RV-C03		No ^b	Not done	Not done
Replace 3',4'-methylenedioxy by methoxy and shorten connecting chain by 2 C-atoms (1 double bond)				






Test CPD.	Change with Respect to Piperine	Active at 10 μ M?	Active at higher conc. (μ M)?	More active than piperine at higher conc. (μ M)?
RV-G01	6'-MeO	No	100 ^e	No
RV-G02	3'-MeO	No	100 ^e	No
RV-G03	4'-MeO	No	100	No
RV-G04	3',4'-Di-MeO	No	100	No

Footnotes

- a. In dose response test.
- b. But the cells displayed weak dendricity suggestive of activity.
- 5 c. This result is considered anomalous. It is intended to re-run the experiment.
- d. Expected to show activity at 50 μ M.
- e. Not corroborated by the t-test at 100 μ M.

Table 7

Code N ^o	Variation on Nitrogen Substituent of Piperine Structure	Effect on melan-a cells at μM concentration			Dendricity
		Percentage cell growth (Repeated experiments) Test cpd.	Stimulant activity	Relative activity to piperine	
RV-A01		183±34** 202±84*	Positive	1.03 1.01	+++
RV-A02		156±58 187±40** 153±19**	Positive	0.5 1.02 0.9	+++
RV-A04		149±47 119±27 147±22	Non-significant here, but positive in dose response test	0.7 0.27 0.66	+
RV-A05		166±35 140±17*	Positive	0.93 0.57	+++

RV-A06		147±66 158±24** 156±40**	170±39** 169±29** 155±18**	Positive	1.69 0.83 1.0	+++
RV-A07		170±24*	216±33*	Positive	0.6	++
RV-A08		200±14	236±17	Positive	0.73	+++
RV-A09		224±19	263±16**	Positive	0.76	+++
RV-A10		308±29**	302±17**	Positive	1.02	+++









RV-A11		264±21**	347±14**	Positive	0.66	+++
*P<0.05, **P<0.01 compared to vehicle treatment (Dunnnett's t test) +++ highly dendritic, ++ moderately dendritic, + weakly dendritic, - no effect						

Table 8

Variation in connecting chain length and amide group		Effect on melan-a cells at μM concentration				
Code N ^o	Structure	Percentage cell growth (Repeated experiments) Test	Piperine	Stimulant activity	Relative activity to piperine	Dendricity
RV-B01		171±33** 148±20 152±22**	180±50** 191±63** 155±18**	Positive	0.88 0.52 0.97	++
RV-B02		140±14 154±33 135±4	180±50** 191±63** 155±18**	Non-significant	0.2 0.59 0.63	+
RV-B03		103±12 116±17	210±65** 170±22**	None	0.02 0.22	-

**P<0.01 compared to vehicle treatment (Dunnett's t test) ++ moderately dendritic, + weakly dendritic, - no effect

Table 9

Code N°	Replacement of amide by ester group Structure	Effect on melan-a cells at μM concentration				
		Percentage cell growth (Repeated experiments) Test	Piperine	Stimulant activity	Relative activity to piperine	Dendricity
RV-AB1		163±38 141±18* 151±7*	210±65** 170±22** 155±18**	Positive	0.57 0.59 0.93	++
RV-AB2		29±9 22±0.4	171±39** 171±39**	Positive	-1 -1.09	Toxic
RV-AB4		224±12**	255±15**	Positive	0.8	++
RV-AB5		166±35**	169±29**	Positive	0.95	++





RV-AB6		148±18**	181±11**	Positive	0.59	+
*P<0.05, **P<0.01 compared to vehicle treatment (Dunnett's t test) +++ highly dendritic, ++ moderately dendritic, + weakly dendritic, - no effect						

Table 10

Replacement of amide by ester group and variation in connecting chain length		Effect on melan-a cells at μM concentration			
Code N ^o	Structure	Percentage cell growth (Repeated experiments)	Stimulant activity	Relative activity to piperine	Dendricity
RV-BB1		Test 149±27 129±15 121±12	210±65** 170±22** 155±18**	Non-significant 0.44 0.41 0.39	+




**P<0.01 compared to vehicle treatment (Dunnett's t test), ++moderately dendricity


Table 11

Reduction of double bonds in connecting chain and variation in chain length		Effect on melan-a cells at μM concentration				
Code N°	Structure	Percentage cell growth (Repeated experiments) Test	Piperine	Stimulant activity	Relative activity to piperine	Dendricity
RV-C02		169±29** 195±89**	180±50** 191±63**	Positive	0.8 1.04	+++
RV-C03		104±5 113±2	171±7** 171±7**	None	0.056 0.18	+


**P<0.01 compared to vehicle treatment (Dunnett's t test) +++ highly dendritic, + weakly dendritic, - no effect

Table 12

Variation in the phenyl substituent and connecting chain length		Effect on melan-a cells at μM concentration				
Code N ^o	Structure	Percentage cell growth (Repeated experiments) Test	Stimulant activity	Relative activity to piperine	Dendricity	
RV-G01		105 \pm 8	None	0.04		
RV-G02		119 \pm 18 87 \pm 17	Negative	0.26 -0.18	-	
RV-G03		121 \pm 8 122 \pm 8*	Non-significant	0.29 0.30	-	

RV-G04		100±9	224±11**	None	0	-
*P<0.05, **P<0.01 compared to vehicle treatment (Dunnnett's t test) - no effect						

Dose response experiments

Code N°	Structure
RV-A01	


Compounds Tested	1μM	10μM	25μM	50μM
Piperine	151±7**♦	202±12**♦	171±15**♦	142±9**
RV-A01	109±7	122±7	142±21**	186±14**□

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5 * P<0.05 Compared to vehicle treatment (Dunnet's t test)

♦ Piperine is significantly more active than test compound P<0.05


□ Test compound is significantly more active than Piperine P<0.05

Code N°	Structure
RV-A02	

Compounds Tested	1µM	10µM	25µM	50µM
Piperine	147±11**♦	192±13**♦	167±19**	142±15**
RV-A02	125±10	167±17**	171±8**	168±12**□

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

- 5 ♦ Piperine is significantly more active than test compound P<0.05
- Test compound is significantly more active than Piperine P<0.05

Code N°	Structure
RV-A04	

Compounds Tested	1μM	10μM	50μM	100μM
Piperine	120±11	178±11**♦	116±13	92±9
RV-A04	101±12	138±10**	150±15**□	71±9
Dentricity RV-A04	-	+	+	

** P<0.01 Compared to vehicle treatment (Dunnet's t test)


5

* P<0.05 Compared to vehicle treatment (Dunnet's t test)

♦ Piperine is significantly more active than test compound P<0.05

□ Test compound is significantly more active than Piperine P<0.05


- no effect, + weakly dendritic, ++ moderately dendritic

Code N°	Structure
RV-A05	

Compounds Tested	1µM	10µM	25µM	50µM
Piperine	173±6**♦	230±13**♦	188±19**	182±15**
RV-A05	155±9**	188±13**	178±18**	174±8**

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5 ♦ Piperine is significantly more active than test compound P<0.05

Code N°	Structure
RV-A06	


Compounds Tested	1 μ M	10 μ M	25 μ M	50 μ M
Piperine	147 \pm 8 \blacklozenge	195 \pm 22**	173 \pm 17*	159 \pm 14
RV-A06	134 \pm 7	188 \pm 14**	172 \pm 15*	135 \pm 24

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5

* P<0.05 Compared to vehicle treatment (Dunnet's t test)

\blacklozenge Piperine is significantly more active than test compound P<0.05

Code N°	Structure
RV-A07	

Compounds Tested	1μM	10μM	50μM	100μM
Piperine	211±16**♦	216±33**	52±15	16±3
RV-A07	140±12**	170±24**	71±5	46±2
Dentricity of RV-A07	++	++	+	+

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5 ♦ Piperine is significantly more active than test compound P<0.05

++ moderately dendritic, + weakly dendritic

Code N°	Structure
RV-A08	


Compounds Tested	1 μ M	10 μ M	50 μ M	100 μ M
Piperine	216 \pm 14** \blacklozenge	236 \pm 17**	61 \pm 11	32 \pm 5
RV-A08	139 \pm 27**	200 \pm 14**	81 \pm 12	62 \pm 13
Dendricity of RV-A08	++	+++	+	+

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5

\blacklozenge Piperine is significantly more active than test compound P<0.05

+++ highly dendritic, ++ moderately dendritic, + weakly dendritic


Code N°	Structure
RV-A09	

Compounds Tested	1 μ M	10 μ M	50 μ M	100 μ M
Piperine	221 \pm 17** \blacklozenge	263 \pm 16**	77 \pm 12	24 \pm 2
RV-A09	187 \pm 15**	224 \pm 19**	85 \pm 5	42 \pm 6
Dendricity of RV-A09	+++	+++	+	+

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5 \blacklozenge Piperine is significantly more active than test compound P<0.05

+++ highly dendritic, + weakly dendritic

Code N°	Structure
RV-A10	


Compounds Tested	1μM	10μM	50μM	100μM
Piperine	236±30**	302±17**	78±11	21±4
RV-A10	301±20**□	308±29**	155±22**□	100±13
Dendricity of RV-A10	+++	+++	++	+

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5

□ Compound is significantly more active than piperine P<0.05

+++ highly dendritic, ++ moderately dendritic, + weakly dendritic

Code N°	Structure
RV-A11	

Compounds Tested	1 μ M	10 μ M	50 μ M	100 μ M
Piperine	251 \pm 19** \blacklozenge	347 \pm 14** \blacklozenge	61 \pm 7	25 \pm 2
RV-A11	189 \pm 6**	264 \pm 21**	158 \pm 20** \square	84 \pm 6
Dendricity of RV-A11	+++	+++	++	+

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5 \blacklozenge Piperine is significantly more active than test compound P<0.05

\square Compound is significantly more active than Piperine P<0.05

Code N°	Structure
RV-B01	


Compounds Tested	1 μ M	10 μ M	25 μ M	50 μ M
Piperine	144 \pm 27** \blacklozenge	190 \pm 7**	172 \pm 11**	153 \pm 10**
RV-B01	111 \pm 6	147 \pm 7**	187 \pm 18**	187 \pm 8** \square

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5

\blacklozenge Piperine is significantly more active than test compound P<0.05

\square Compound is significantly more active than Piperine P<0.05

Code N°	Structure
RV-AB1	

Compounds Tested	1μM	10μM	50μM	100μM
Piperine	133±31**	177±14***◆	139±16*	95±24
RV-AB1	125±13	147±16**	187±12***□	171±8***□
Dendricity of RV-AB1	-	+	++	++

** P<0.01 Compared to vehicle treatment (Dunnet's t test)


5

* P<0.05 compared to vehicle treatment (Dunnett's t test)

◆ Piperine is significantly more active than test compound P<0.05

□ Compound is significantly more active than Piperine P<0.05

- no effect, + weakly dendritic, ++ moderately dendritic

Code N°	Structure
RV-AB4	

Compounds Tested	1 μ M	10 μ M	50 μ M	100 μ M
Piperine	223 \pm 18** \blacklozenge	255 \pm 15**	60 \pm 16	24 \pm 6
RV-AB4	175 \pm 6**	224 \pm 12**	-148 \pm 19** \square	90 \pm 7
Dendricity of RV-AB4	++	++	++	+


** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5

\blacklozenge Piperine is significantly more active than test compound P<0.05

\square Compound is significantly more active than Piperine P<0.05

++ moderately dendritic, + weakly dendritic

Code N°	Structure
RV-AB5	

Compounds Tested	1μM	10μM	50μM	100μM
Piperine	141±26**◆	220±29**◆	45±12	23±4
RV-AB5	120±21	151±19**	163±8**□	123±8
Dendricity of RV-AB5	-	++	++	+


** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5

◆ Piperine is significantly more active than test compound P<0.05

□ Compound is significantly more active than Piperine P<0.05

++ moderately dendritic, + weakly dendritic

Code N°	Structure
RV-AB6	


Compounds Tested	1 μ M	10 μ M	50 μ M	100 μ M
Piperine	113 \pm 10	181 \pm 11**	43 \pm 6	23 \pm 6
RV-AB6	103 \pm 5	148 \pm 18**	190 \pm 11** <input type="checkbox"/>	128 \pm 17** <input type="checkbox"/>
Dendricity of RV-AB6	-	+	++	+

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5

Compound is significantly more active than Piperine P<0.05


++ moderately dendritic, + weakly dendritic, - no effect

Code N°	Structure
RV-C02	

Compounds Tested	1μM	10μM	25μM	50μM
Piperine	158±10**♦	203±11**	188±12**	164±6**
RV-C02	134±15**	183±33**	199±31**	175±12**

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5 ♦ Piperine is significantly more active than test compound P<0.05


Code N°	Structure
RV-G01	

Compounds Tested	1 μ M	10 μ M	50 μ M	100 μ M
Piperine	161 \pm 23**	202 \pm 29**	61 \pm 5	40 \pm 7
RV-G01-	99 \pm 8	105 \pm 8	103 \pm 6	119 \pm 9
Dendricity of RV-G01	-	-	-	-

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5

- no effect


Code N°	Structure
RV-G02	

Compounds Tested	1µM	10µM	50µM	100µM
Piperine	151±17**	201±15**	57±15	39±11
RV-G02	99±5	95±18	110±11	127±9
Dendricity of RV-G02	-	-	-	-

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5

- no effect


Code N°	Structure
RV-G03	

Compounds Tested	1μM	10μM	50μM	100μM
Piperine	163±9**	181±23**	59±11	40±12
RV-G03	90±10	108±20	111±10	133±15**
Dendricity of RV-G03	-	-	-	-

** P<0.01 Compared to vehicle treatment (Dunnet's t test)

5

- no effect

Code N°	Structure
RV-G04	

Compounds Tested	1 μ M	10 μ M	50 μ M	100 μ M
Piperine	179 \pm 12**	224 \pm 11**	92 \pm 19	35 \pm 4
RV-G04	95 \pm 11	100 \pm 9	114 \pm 8	123 \pm 7*
Dendricity of RV-G04	-	-	-	-

** P<0.01 Compared to vehicle treatment (Dunnet's t test)


5

- no effect

2. Synthesis of amide derivatives of piperinic acid

2.1 Preparation of piperinic acid (RV-A00)

To piperine (1) (2g, 0.7mmol, 1eq), 20% of methanolic KOH (100ml) was added and refluxed for 2days. After completion of the hydrolysis, methanol was removed under reduced pressure and a yellow coloured oily solid was obtained. This residue was dissolved in water (50ml) and acidified with 6N HCl to pH <1 yielding a yellowish precipitate of piperinic acid. Recrystallization from methanol gave yellow needles (0.9g, 60% yield). m.p. 206^o-208^oC (Lit m.p. 217^o-218^oC)¹




10

2.2 Synthesis of piperlonguminine (RV-A06)

A mixture of piperinic acid (350mg, 0.0016mole, 1eq) and triethylamine (0.4ml, 0.0032mole, 2eq) in dichloromethane (50ml) was stirred for 15min at 0^oC. To this mixture methanesulfonyl chloride (0.18ml, 0.0024mole, 1.5eq) was added and stirred for further 30 min at 0^oC. Isobutylamine (0.23ml, 0.0024mole, 1.5eq) was added to the mixture and stirred for 1h at 0^oC and 2h at room temperature. Dichloromethane (50ml) was added to the mixture which was then washed with 5% HCl (3x100ml), saturated aqueous NaHCO₃ (3x100ml) and water (3x100ml). The organic fraction was dried over anhydrous sodium sulphate, filtered and rotary evaporated to yield a yellowish solid residue. Recrystallisation from methanol yielded colourless needles of piperlonguminine (120mg, 32% yield)². The reaction is presumed to proceed through a mesylate ester intermediate.

20

Piperlonguminine (RV-A06)



$^1\text{H-NMR}$ (CDCl_3) δ : 5.96 (d, 1H, $J=14.8$, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 7.36 (d,d, 1H, $J=10.5$, 14.8, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.66 (d,d, 1H, $J=15.4$, 10.5, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.76 (d, 1H, $J=15.4$, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.96 (d, 1H, $J=1.6$, Ar-7H), 6.76 (d, 1H, $J=8.0$, Ar-10H), 6.87 (d, d, 1H, $J=1.6$, 8.0 Ar-11H), 5.97 (s, 2H, O-CH₂-O), 3.18 (t, 2H, $J=6.5$ CH₂-CH), 1.83 (m, 1H, $J=6.5$ CH₂-CH), 0.94 (d, 6H, $J=6.5$, (CH₃)₂), \rightarrow

5.82 (t, 1H, NH $J=5.3$)

$^{13}\text{C-NMR}$ (CDCl_3) : 20.4 (CH₃), 29.4 (CH), 47.3 (CH₂), 102.2 (CH₂), 106.2 (CH), 109.1 (CH), 123.3 (CH), 125.5 (CH), 126.0 (CH), 132.0 (C), 138.0 (CH), 140.4 (CH), 148.9 (C), 149.2 (C), 166.2 (C)

MS m/z (%): 273 (M^+ 98), 216 (20), 201 (100), 174 (25), 173 (65), 172 (23), 171 (17), 143 (20), 115 (40), 96 (11).

IR (KBr) : ν_{max} (carbonyl group) 1644

m.p. 161.2^o-161.7^oC (Lit m.p. 156^o-160^oC)¹


15

2.3 Synthesis of other amide derivatives of piperinic acid

The general method was as for piperlonguminine (section 1.2), using the same proportions of reactive amine, triethylamine and methanesulfonyl chloride relative to piperinic acid (200 or 300 mg., 1eq). Recrystallisation from ethyl acetate/petroleum spirit yielded the other amide derivatives of piperinic acid.

20

5-E,E-piperinoylpyrrolidine (RV-A01)



¹H-NMR (CDCl₃) δ: 6.26 (d, 1H, J=14.7, CH=CH-CH=CH), 7.43 (d,d, 1H, J=9.5, 14.7, CH=CH-CH=CH), 6.73 (d,d, 1H, J=15.3, 9.5, CH=CH-CH=CH), 6.78 (d, 1H, J=15.3 CH=CH-CH=CH), 6.98 (d,1H J=1.6, Ar-7-H), 6.77 (d,1H J=8.0, Ar-10-H), 6.89 (d, d, 1H J=1.6, 8.0 Ar-11-H), 5.97 (s, 2H, O-CH₂-O), 3.57 (t, 2H, J=4.0 N- CH₂ (pyrrolidine)) 3.54 (t, 2H, J=4.0 N- CH₂ (pyrrolidine)) 1.90 (m, 2H, CH₂.CH₂(pyrrolidine)) 1.87 (m, 2H, CH₂.CH₂(pyrrolidine))

¹³C-NMR (CDCl₃) : 24.3 (CH₂), 26.1 (CH₂), 45.9 (CH₂), 46.4 (CH₂), 101.2 (CH₂), 105.7 (CH), 108.4 (CH), 121.4 (CH), 122.5 (CH), 125.2 (CH), 130.9 (C), 138.7 (CH), 141.7 (CH), 148.1 (C), 148.2 (C), 164.9 (C)


MS m/z (%) : 271 (M⁺ 78), 201 (100), 173 (30), 172 (15), 171 (13) 143 (13), 115 (27)

IR (KBr) : ν_{max} (carbonyl group) 1637

m.p. 142.9^o-143^oC (Lit m.p. 142^o-143^oC)², yield 49.2%

15

5-E,E-piperinoyl morpholine (RV-A02)



¹H-NMR (CDCl₃) δ: 6.37 (d, 1H, J=14.6, CH=CH-CH=CH), 7.45 (d,d, 1H, J=10.2, 14.6, CH=CH-CH=CH), 6.72 (d,d, 1H, J=15.5, 10.2, CH=CH-CH=CH), 6.79 (d, 1H, J=15.5 CH=CH-CH=CH), 6.98 (d,1H J=1.5, Ar-7-H), 6.80 (d,1H J=8.0, Ar-10-H), 6.89 (d, d, 1H J=1.5, 8.0 Ar-11-H), 5.98 (s, 2H, O-CH₂-O), 3.70 (t, 2H, J=4.0 CH₂-N- CH₂ (morpholine)) 3.60 (t, 2H, J=4.0 CH₂-O- CH₂ (morpholine))


$^{13}\text{C-NMR}$ (CDCl_3) : 42.3 (CH_2), 46.1(CH_2), 66(CH_2), 66(CH_2), 101.3 (CH_2), 106.5 (CH), 108.5 (CH), 118.7 (CH), 122.7 (CH), 124.9 (CH), 130.8 (C), 139.1 (CH), 143.4 (CH), 148.2 (C), 148.3 (C), 165.6 (C)

MS m/z (%) : 287 (M^+ 57), 201 (100), 173 (25), 171 (10) 143 (10), 115 (30)

5 IR (KBr): ν_{max} (carbonyl group) 1641

m.p. 161.8 $^{\circ}$ -162.5 $^{\circ}$ C (Lit m.p. 167-168 $^{\circ}$ C)³, yield 44.1%

5-E,E-piperinoylpiperinolymine (RV-A04)




10 $^1\text{H-NMR}$ (CDCl_3) δ : 5.98(d, 1H, J=14.9, CH=CH-CH=CH), 7.34 (d,d, 1H, J=10.7, 14.9, CH=CH-CH=CH), 6.73 (d,d, 1H, J=15.5, 10.7, CH=CH-CH=CH), 6.79 (d, 1H, J=15.5 CH=CH-CH=CH), 6.98 (d,2H J=1.5, Ar-7,3'-H), 6.78 (d,2H J=8.0, Ar-10,6'-H), 6.89 (d, d, 2H J=1.6, 8.0 Ar-11,7'-H), 5.98 (s, 2H, O-CH₂-O), 5.93 (s, 2H, O-CH₂-O), 4.40 (d, 2H, CH₂) 3.57 (br, 1H, NH)

15 $^{13}\text{C-NMR}$ (CDCl_3) : 43.4 (CH_2), 101.1 (CH_2), 101.4 (CH_2), 105.8 (CH), 108.3(CH) 108.5 (CH), 108.6 (CH), 121.2 (CH), 122.8 (CH), 124.7 (CH), 130.9 (C), 132.2 (C) 139.9 (CH), 141.6 (CH), 147.0 (C) 147.9 (C)148.3 (C), 148.4(C), 166.9 (C)

MS m/z (%) : 351 (M^+ 81), 216 (15), 203 (12), 202 (53) 201 (29), 174 (31), 173 (22), 150 (23) 144 (11), 143 (10), 135 (100), 116 (12) 115(29)


20 m.p. 190.5 $^{\circ}$ -191.7 $^{\circ}$ C, yield 50.1%

5-E,E-piperinoylhexylamine RV-A05




¹H-NMR (CDCl₃) δ: 5.90 (d, 1H, J=14.8, CH=CH-CH=CH), 7.35 (d,d, 1H, J=10.6, 14.8, CH=CH-CH=CH), 6.66 (d,d, 1H, J=15.4, 10.6, CH=CH-CH=CH), 6.76 (d, 1H, J=15.4 CH=CH-CH=CH), 6.97 (d,1H J=1.4, Ar-7H), 6.77 (d,1H J=8.0, Ar-10H), 6.88 (d, d, 1H J=1.5, 8.0 Ar-11H), 5.97 (s, 2H, O-CH₂-O), 3.34 (q, 2H, CH₂ - CH₂- CH₂- CH₂- CH₂)
 5 1.54 (m, 2H, CH₂ - CH₂- CH₂- CH₂- CH₂) 1.32 (m, 6H, CH₂ - CH₂- CH₂- CH₂- CH₂)
 0.88 (t, 3H, CH₃), 5.54 (br, NH)
¹³C-NMR (CDCl₃) : 14.3 (CH₃), 22.5 (CH₂), 26.6 (CH₂), 29.6 (CH₂), 31.5 (CH₂), 39.7 (CH₂), 101.3 (CH₂), 105.7 (CH), 108.5 (CH), 122.5 (CH), 123.2 (CH), 124.6 (CH), 130.8 (C), 138.7 (CH), 140.9 (CH) 148.2 (C), 148.2 (C), 166.0 (C)
 10 MS m/z (%) : 301(M⁺94), 202 (18) 201 (73), 174 (40), 173 (100), 172 (31), 171 (15) 143 (24), 115 (63)
 IR (KBr) : ν_{max} (carbonyl group)1641
 m.p. 149.5⁰-149.8⁰C (Lit m.p. 139⁰-141⁰C)⁴, yield 50.1%

15 **5-E,E-piperinoylmethylamine (RV-A07)**



¹H-NMR (CDCl₃) δ: 5.91 (d, 1H, J=14.8, CH=CH-CH=CH), 7.36 (d,d, 1H, J=10.7, 14.8, CH=CH-CH=CH), 6.66 (d,d, 1H, J=15.4, 10.6, CH=CH-CH=CH), 6.77 (d, 1H, J=15.4 CH=CH-CH=CH), 6.97 (d,1H J=1.5, Ar-7H), 6.77 (d,1H J=8.0, Ar-10H), 6.88 (d, d, 1H J=1.6, 8.0 Ar-11H), 5.97 (s, 2H, O-CH₂-O), 2.91(t, 3H, CH₃), 5.61 (br, NH)
 20 ¹³C-NMR (CDCl₃) : 26.9 (CH₃), 101.7 (CH₂), 106.1 (CH), 108.9 (CH), 123.0 (CH), 123.3 (CH), 125.0 (CH), 131.2 (C), 139.2 (CH), 141.4 (CH), 148.6 (C), 148.6 (C), 167.2 (C)
 MS m/z (%) : 231(M⁺89), 201 (42), 173 (67), 172 (32), 171 (17), 143 (27), 116 (21) 115 (100), 89 (12)
 25 m.p. 181.1⁰-182.4⁰C (Lit m.p. 186⁰C)⁵, yield 48.2%

5-E,E-piperinoylethylamine (RV-A08)




$^1\text{H-NMR}$ (CD_3OD) δ : 6.14 (d, 1H, $J=15.0$, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 7.37 (d,d, 1H, $J=10.2$,
15.0, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.93 (d,d, 1H, $J=15.7$, 10.6, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.87 (d, 1H,
5 $J=15.7$ $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.97 (d, 1H $J=1.5$, Ar-7H), 6.77 (d, 1H $J=8.0$, Ar-10H), 6.88 (d,
d, 1H $J=1.6$, 8.0 Ar-11H), 5.97 (s, 2H, O- CH_2 -O), 3.39 (m, 2H, $J=6.2$, CH_2), 1.22 (t, 3H,
 $J=6.1$, CH_3),

$^{13}\text{C-NMR}$ (CDCl_3) : 14.7 (CH_3), 36.9 (CH_2), 103.2 (CH_2), 107.2 (CH), 109.8 (CH),
121.2 (CH), 124.9 (CH), 125.9 (CH), 132.4 (C), 142.9 (CH), 145.2 (CH), 150.2 (C),
10 150.6 (C), 170 (C)

MS m/z (%) : 245 (M^+ 78), 218 (34), 201 (71), 200 (49), 174 (64), 173 (80), 172 (76), 171
(65), 143 (75), 116 (68), 115 (100)

m.p. 158.5 $^{\circ}$ -159.9 $^{\circ}$ C (Lit m.p. 162 $^{\circ}$ -164 $^{\circ}$ C)⁴, yield 45.6%

5-E,E-piperinoylisopropylamine (RV-A09)




¹H-NMR (CDCl₃) δ: 5.87 (d, 1H, J=14.8, CH=CH-CH=CH), 7.36 (d,d, 1H, J=10.7, 14.8, CH=CH-CH=CH), 6.66 (d,d, 1H, J=15.4, 10.6, CH=CH-CH=CH), 6.76 (d, 1H, J=15.2 CH=CH-CH=CH), 6.97 (d,1H J=1.6, Ar-7H), 6.77 (d,1H J=8.0, Ar-10H), 6.88 (d,d, 1H J=1.6, 8.0 Ar-11H), 5.97 (s, 2H, O-CH₂-O), 4.15(m, 1H, J=6.6, CH), 5.36 (d, 1H, J=7.3 NH), 1.19 (d, 6H, J=6.6, (CH₃)₂)

¹³C-NMR (CDCl₃) : 23.2 (CH₃)₂, 41.9 (CH), 101.9 (CH₂), 106.4 (CH), 108.9 (CH), 123.0 (CH), 123.8 (CH), 124.1 (CH), 131.3 (C), 140.2 (CH), 141.2 (CH), 148.8 (C), 148.6 (C) 165.6 (C)

MS m/z (%) : 259(M⁺80), 201 (62), 174 (34), 173 (74), 172 (31), 171 (15), 143 (30), 116 (16), 115 (100)

m.p. 169⁰-169.4⁰C (Lit m.p. 171⁰-173⁰C)⁴, yield 52%

5-E,E-piperinoyl cyclohexylamine (RV-A10)




¹H-NMR (CDCl₃) δ: 5.93 (d, 1H, J=14.8, CH=CH-CH=CH), 7.35 (d,d, 1H, J=10.6, 14.8, CH=CH-CH=CH), 6.66 (d,d, 1H, J=15.3, 10.6, CH=CH-CH=CH), 6.76 (d, 1H, J=15.4 CH=CH-CH=CH), 6.96 (d,1H J=1.6, Ar-7H), 6.76 (d,1H J=8.0, Ar-10H), 6.87 (d, d, 1H J=1.6, 8.0 Ar-11H), 5.97 (s, 2H, O-CH₂-O), 3.87 (m, 1H, CH (cyclohexyl)) 1.99 (m, 2H, CH₂(cyclohexyl)) 1.65 (m, 4H, CH₂-CH₂(cyclohexyl)) 1.39 (m, 2H, CH₂(cyclohexyl)) 1.18 (m, 2H, CH₂(cyclohexyl)) 5.48 (d,J=8.0 NH)

^{13}C -NMR (CDCl_3) : 25.3 ($(\text{CH}_2)_2$), 25.9 (CH_2), 33.6 ($(\text{CH}_2)_2$), 48.6 (CH), 101.3 (CH_2), 101.7 (CH), 106.1 (CH), 108.9 (CH), 123.0 (CH), 124.0 (CH), 125.1 (CH), 131.3 (C), 139.0 (CH), 141.2 (CH) 148.5 (C), 148.5 (C), 165.5 (C)

MS m/z (%) : 299(M^+56), 259 (48) 216 (33), 201 (60), 174 (33), 173 (61), 172 (18), 171 (16) 143 (17), 115 (100)

m.p. 196.4^0 - 197.3^0C (Lit m.p. 199^0 - 200^0C)⁴, yield 57.4%

5-E,E-piperinoylbutylamine (RV-A11)



10

^1H -NMR (CDCl_3) δ : 5.97 (d, 1H, $J=14.8$, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 7.35 (d,d, 1H, $J=10.7$, 14.8, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.66 (d,d, 1H, $J=15.4$, 10.6, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.76 (d, 1H, $J=15.4$, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.97 (d, 1H $J=1.6$, Ar-7H), 6.77 (d, 1H $J=8.0$, Ar-10H), 6.89 (d, d, 1H $J=1.5$, 8.0 Ar-11H), 5.97 (s, 2H, O- CH_2 -O), 3.36 (q, 2H, CH_2 - CH_2 - CH_2 -) 1.54 (m, 2H,

15

CH_2 - CH_2 - CH_2) 1.39 (m, 6H, CH_2 - CH_2 - CH_2) 0.93 (t, 3H, CH_3), 5.47 (br, NH)

^{13}C -NMR (CDCl_3) : 14.2 (CH_3), 20.5 (CH_2), 32.2 (CH_2), 39.8 (CH_2), 101.7 (CH_2), 106.1 (CH), 108.9 (CH), 123.0 (CH), 123.6 (CH), 125.0 (CH), 131.3 (C), 139.2 (CH), 141.3 (CH) 148.6 (C), 148.6 (C), 166.4 (C)

m.p. 144.2^0 - 145.6^0C (Lit m.p. 144^0 - 145^0C)⁴, yield 38.4%

20

References:

¹Chatterjee, A., and Dutta, C.P. (1967). Alkaloids of *Piper longum* Linn-I Structure and synthesis of piperlongumine and piperlonguminine, *Tetrahedron*, **23**, 1769-1781.

25 ²Nokio Nakumara, Fumiyuki Kiuchi, and Yoshisuke Tsuda (1988). Infrared spectra of conjugated amides: Reassignment of the C=O and C=C absorptions: *Chemical and Pharmaceutical Bulletin*, **36**, 2647-2651.

³H.Oediger and A.Schulze (Bayer AG), (1979), *Deutsche Auslegeschrift* 2757 483

⁴Paula, Vanderlucia F. de; A Barbosa, Luiz C. de; Demuner, Antonio J.; Pilo-Veloso, Dorila; Picanco, Marcelo C. (2000) Pest Management Science **56**, 2, 168 – 174.

⁵Gokale et al., (1948) Journal of University Bombay Science **16/5A** 32-35

5 **3. Synthesis of ester derivatives of piperinic acid**

3.1 Preparation of piperinic acid (RV-A00)

As described above.

3.2 Synthesis of 5-E,E-piperinic acid methyl ester (RV-AB1)


10 A mixture of piperinic acid (300mg, 0.0014mole, 1eq) and triethylamine (0.39ml, 0.0028mole, 2eq) in dichloromethane (50ml) was stirred for 15min at 0°C. To this mixture methanesulfonyl chloride (0.16ml, 0.0021mole, 1.5eq) was added and stirred for further 30 min at 0°C. Methanol in excess (10ml) was added to the mixture and stirred for 15 1h at 0°C and 1h at room temperature. Dichloromethane (50ml) was added to the mixture which was then washed with water (3x100ml), 5% NaHCO₃ (3x100ml) and water (3x100ml). The organic fraction was dried over anhydrous sodium sulphate, filtered and rotary evaporated to yield a yellowish solid residue. Recrystallisation from ethyl acetate/petroleum spirit yielded ester (180mg, 56.2% yield). m.p. 142.9⁰-143⁰C (Lit m.p. 140⁰C)⁶

20

3.3 Synthesis of other esters of piperinic acid.

They were synthesised as described in section 3.2, replacing methanol (10ml) ethanol (10ml), isopropanol, butanol or propanol (15ml).

25 **5-E,E-piperinic acid methyl ester (RV-AB1)**




$^1\text{H-NMR}$ (CDCl_3) δ : 5.94 (d, 1H, $J=15.2$, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 7.41 (d,d, 1H, $J=10.8$, 15.2, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.70 (d,d, 1H, $J=15.4$, 10.8, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.81 (d, 1H, $J=15.7$, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.99 (d, 1H $J=1.6$, Ar-7H), 6.79 (d, 1H $J=8.1$, Ar-10H), 6.91 (d, d, 1H $J=1.5$, 8.1 Ar-11H), 5.98 (s, 2H, $\text{O}-\text{CH}_2-\text{O}$), 3.57 (t, 3H, br, OCH_3 $J=4.7$)

5 $^{13}\text{C-NMR}$ (CDCl_3) δ : 51.5 (CH_3), 101.8 (CH_2), 106.2 (CH) 108.9 (CH), 120.0 (CH), 123.4 (CH) 124.7 (CH), 130.8 (CH), 140.9 (C), 145.5 (CH), 148.6 (C), 148.9 (C), 168.9 (C)

MS m/z (%) : 232 (M^+ 69), 201 (19), 174 (12), 173 (100), 172 (39), 171 (12) 143 (33), 116 (11), 115 (53) 101 (15), 100 (12)

10

5-E,E-piperinic acid ethyl ester (RV-AB2)




15 $^1\text{H-NMR}$ (CDCl_3) δ : 5.94 (d, 1H, $J=15.2$, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 7.41 (d,d, 1H, $J=10.8$, 15.3, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.70 (d,d, 1H, $J=15.4$, 10.8, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.81 (d, 1H, $J=15.5$, $\text{CH}=\text{CH}-\text{CH}=\text{CH}$), 6.99 (d, 1H $J=1.6$, Ar-7H), 6.78 (d, 1H $J=8.1$, Ar-10H), 6.91 (d, d, 1H $J=1.6$, 8.1 Ar-11H), 5.98 (s, 2H, $\text{O}-\text{CH}_2-\text{O}$), 4.22 (q, 2H, OCH_2 $J=7.2$), 1.31 (t, 3H, CH_3 $J=7.2$)

20 $^{13}\text{C-NMR}$ (CDCl_3) : 14.7 (CH_3), 60.7 (CH_2), 101.6 (CH_2), 106.3 (CH) 108.9 (CH), 120.8 (CH), 123.3 (CH) 124.9 (CH), 131.0 (CH), 140.5 (CH), 145.1 (CH), 148.7 (C), 148.9 (C), 167.6 (C)

5-E,E-piperinic acid isopropyl ester (RV-AB4)

Physical data are not available for this compound.

5-E,E-piperinic acid propyl ester (RV-AB5)



¹H-NMR (CDCl₃) δ: 5.94 (d, 1H, J=15.2, CH=CH-CH=CH), 7.41 (d,d, 1H, J=10.7, 15.2, CH=CH-CH=CH), 6.70 (d,d, 1H, J=15.4, 10.8, CH=CH-CH=CH), 6.76 (d, 1H, J=15.4 CH=CH-CH=CH), 6.99 (d,1H J=1.6, Ar-7H), 6.78 (d,1H J=8.1, Ar-10H), 6.91 (d, d, 1H J=1.5, 8.0 Ar-11H), 5.98 (s, 2H, O-CH₂-O), 4.12 (t, 2H, OCH₂ J=6.7) 1.69 (m, 2H, CH₂ J=7.3) 0.97 (t, 3H, CH₃ J=7.4)

¹³C-NMR (CDCl₃) :10.9 (CH₃), 22.5 (CH₂), 66.3 (CH₂),101.8 (CH₂), 106.2 (CH) 108.9 (CH), 120.9 (CH), 123.3 (CH)124.9 (CH), 131.0 (CH), 140.5 (CH), 145.1 (CH), 148.7 (C), 148.9(C), 167.7 (C)


MS m/z (%) : 260 (M⁺ 59), 201 (26), 174 (18), 173 (100), 172 (39), 171 (14) 143 (34), 116 (16), 115 (73),100 (12)

m.p. 119^o-120^oC

15 **References**

⁶Avijit Banerjee, Tapasree Ghosal, and Aditi Kacharya. (1984). Indian Journal of Chemistry, 23B, 546-549.

5-E,E-piperinic acid butyl ester (RV-AB6)



20 ¹H-NMR (CDCl₃) δ: 5.94 (d, 1H, J=15.2, CH=CH-CH=CH), 7.40 (d,d, 1H, J=10.7, 15.3, CH=CH-CH=CH), 6.70 (d,d, 1H, J=15.4, 10.8, CH=CH-CH=CH), 6.76 (d, 1H, J=15.4 CH=CH-CH=CH), 6.99 (d,1H J=1.6, Ar-7H), 6.78 (d,1H J=8.0, Ar-10H), 6.91 (d, d, 1H

J=1.5, 8.0 Ar-11H), 5.98 (s, 2H, O-CH₂-O), 4.12 (t, 2H, OCH₂ J=6.7) 1.69 (m, 2H, CH₂ J=7.3) 1.69 (m, 2H, CH₂ J=7.6), 0.95 (t, 3H, CH₃ J=7.5)


MS m/z (%) : 274 (M⁺ 50), 201 (15), 174 (14), 173 (100), 172 (30), 171 (14) 143 (21), 115 (55)

5 Obtained as an oil.

4. Synthesis of amide derivatives of 3,4-methylenedioxcinnamic acid

These 3,4-methylenedioxcinnamide derivatives were synthesised as described in Section 2.2, but using 3,4-methylenedioxcinnamic acid (500mg) as the starting acid and
10 reducing the proportion of triethylamine to 1.5 equivalent with respect to the starting acid. Also, in the first stage, the reaction mixture was stirred for 2 hours, instead of 30 minutes, again at 0°C.

1-(3-trans-benzo-1,3-dioxol-5-ylacryloyl)piperidine (RV-B01)




15

¹H-NMR (CDCl₃) δ: 7.56 (d, 1H, J=15.3, CH=CH), 6.73 (d, 1H, J=15.3, CH=CH-), 7.03 (d, 1H J=1.5, Ar-7H), 6.79 (d, 1H J=8.0, Ar-8H), 6.99 (d, d, 1H J=1.6, 8.0 Ar-9H), 5.98 (s, 2H, O-CH₂-O), 3.57 (br, 2H, CH₂-N-CH₂), 3.65 (br, 2H, CH₂-N-CH₂(piperidine)), 1.65 (m, 6H, CH₂-CH₂-CH₂- (piperidine))

20 ¹³C-NMR (CDCl₃) : 24.8 (CH₂), 25.6 (CH₂), 26.7 (CH₂), 43.3 (CH₂), 46.9 (CH₂), 101.3 (CH₂), 106.7 (CH), 108.4 (CH), 115.6 (CH), 123.5 (CH), 129.9 (C), 141.9 (CH) 148.1 (CH), 148.8 (C), 165.4 (C)


m.p. 80.1^o-82^oC (Lit m.p. 80^o-82^oC)⁷, yield 49.2%

1-(3-trans-benzo-1,3-dioxol-5-ylacryloyl)pyrrolidine (RV-B02)



- 5 $^1\text{H-NMR}$ (CDCl_3) δ : 7.60 (d, 1H, $J=15.2$, $\text{CH}=\text{CH}$), 6.73 (d, 1H, $J=15.3$, $\text{CH}=\text{CH}$), 7.04 (d, 1H $J=1.5$, Ar-7H), 6.80 (d, 1H $J=8.0$, Ar-8H), 7.01 (d, d, 1H $J=1.5$, 8.0 Ar-9H), 5.99 (s, 2H, O- CH_2 -O), 3.61 (br, 2H, CH_2 -N- CH_2 (pyrrolidine)), 3.57 (br, 2H, CH_2 -N- CH_2 (pyrrolidine)), 1.99 (4H, CH_2 - CH_2 (pyrrolidine)),
- $^{13}\text{C-NMR}$ (CDCl_3) : 24.3 (CH_2), 26.1 (CH_2) 46.0 (CH_2), 46.5 (CH_2), 101.4 (CH_2), 106.4 (CH), 108.5 (CH), 116.8 (CH), 123.8 (CH), 129.7 (C), 141.0 (CH) 148.1 (C), 148.9 (C), 164.8 (C)
- 10 MS m/z (%) : 245(M^+ 62), 176 (41) 175 (100) 145 (36), 117 (11), 89 (14).
m.p. 152.5 $^\circ$ -153 $^\circ$ C, yield 44.1%

1-(3-trans-benzo-1,3-dioxol-5-ylacryloyl)morpholine (RV-B03)




- 15 $^1\text{H-NMR}$ (CDCl_3) δ : 7.61 (d, 1H, $J=15.3$, $\text{CH}=\text{CH}$), 6.73 (d, 1H, $J=15.3$, $\text{CH}=\text{CH}$), 7.03 (d, 1H $J=1.4$, Ar-7H), 6.80 (d, 1H $J=8.0$, Ar-8H), 7.01 (d, d, 1H $J=1.4$, 8.0 Ar-9H), 5.99 (s, 2H, O- CH_2 -O), 3.72 (br, 4H, CH_2 -N- CH_2 (morpholine)), 3.67 (br, 4H, CH_2 -O- CH_2 (morpholine)),
- $^{13}\text{C-NMR}$ (CDCl_3) : 42.6 (CH_2), 46.2 (CH_2), 66.8 (CH_2), 46.5 (CH_2), 101.4 (CH_2), 106.3 (CH), 108.5 (CH), 114.4 (CH), 123.9 (CH), 129.5 (CH), 143.0 (CH) 148.2 (C), 148.9 (C), 149.1 (C), 165.6 (C)
- 20 MS m/z (%) : 261(M^+ 60), 176 (24) 175 (100) 145 (30), 117 (10), 89 (11).
m.p. 160 $^\circ$ -160.3 $^\circ$ C, yield 50.1%

5. Synthesis of 3-trans-benzo-1,3-dioxol-5-ylacrylic acid methyl ester (RV-BB1)

To 3,4-methylenedioxcinnamic acid (2g, 0.01mol, 1eq) methanol (4ml, 10eq) was added. Sulphuric acid (0.2ml) was added and refluxed overnight. The solvent was rotary evaporated to yield solid residue. This residue was dissolved in ether and washed with water (2x100ml) and 5%NaHCO₃ (3x100ml) and with water (2x100ml). The organic fraction was dried over anhydrous sodium sulphate and rotary evaporated to yield white solid. Recrystallisation from ethyl acetate/petroleum spirit yielded crystals (69.4% yield) m.p. 133.7^o-134.2^oC (Lit m.p.134^oC)⁸

10

3-trans-benzo-1,3-dioxol-5-ylacrylic acid methyl ester (RV-BB1)



¹H-NMR (CDCl₃) δ: 7.59 (d, 1H, J=15.9, CH=CH), 6.26 (d, 1H, J=15.9, CH=CH-), 7.03 (d, 1H J=1.5, Ar-7H), 6.81 (d, 1H J=8.0, Ar-8H), 7.01 (d, d, 1H J=1.5, 8.0 Ar-9H), 6.00 (s, 2H, O-CH₂-O), 3.79 (s, 3H, OCH₃)

15

¹³C-NMR (CDCl₃) : 51.6 (CH₃), 101.5 (t CH₂), 106.5 (CH), 108.5 (CH), 115.7 (CH), 124.4 (CH), 128.8 (CH), 144.5 (CH) 148.3 (C), 148.6 (C), 148.2 (C), 167.6 (C)

MS m/z (%) : 206(M⁺ 100), 175 (68) 175 (100) 145 (27), 117 (10), 89 (11).

20


References:

⁷H. Staudinger and H.Schneider. (1923). Chem. Ber. 56, 699.

⁸Takemoto *et al.* (1985). Chemical and Pharmaceutical Bulletin 23, 1161.


6. Synthesis of tetrahydropiperine (RV-C02)

Piperine (2g, 7mmol) was hydrogenated in ethanol (50ml) over 5% Pd-C under a pressure of hydrogen at 10 psi for 30mins to give tetrahydropiperine (1.59g, 78%yield) as an oil².



5

Tetrahydropiperine (RV-C02)



¹H-NMR (CDCl₃) δ: 2.55 (t, 4H, J=7.0 CH₂-CH₂-CH₂-CH₂), 2.32 (t, 4H, J=7.0 CH₂-CH₂-CH₂-CH₂) 6.66 (d, 1H J=1.3, Ar-7H), 6.70 (d, 1H J=8.0, Ar-10H), 6.61 (d, d, 1H J=1.2, 8.0 Ar-11H), 5.89 (s, 2H, O-CH₂-O), 3.53 (t, 2H, N-CH₂ (piperidine)) 3.35 (t, 2H, N-CH₂ (piperidine)) 1.63 (m, 2H, CH₂-CH₂-CH₂ (piperidine)) 1.54 (m, 2H, CH₂-CH₂(piperidine))


¹³C-NMR (CDCl₃) : 24.5 (CH₂), 24.9 (CH₂), 25.5 (CH₂), 26.5 (CH₂), 31.4 (CH₂), 33.2 (CH₂), 35.4 (CH₂), 42.5(CH₂), 46.6(CH₂), 100.7 (CH₂), 108.0 (CH), 108.8 (CH), 109.0 (CH), 121.0 (C), 145.4 (C) 147.4 (C), 171.1 (C)

MS m/z (%) : 289 (M⁺ 71), 204 (31), 154 (23), 148 (22), 141 (23), 140 (38), 135 (28) 127 (100), 112 (23), 86 (12), 84 (24), 70 (10), 36 (11)

7. Synthesis of 3-benzo-1,3-dioxol-5-ylpropionic acid piperidide

7.1 Synthesis of 3-benzo-1,3-dioxol-5-ylpropionic acid

3-benzo-1,3-dioxol-5-ylacrylic acid (2g) was hydrogenated in ethanol (50ml) over 5% Pd-C under a pressure of hydrogen at 10 psi for 40mins to give 3-benzo-1,3-dioxol-5-ylpropionic acid (1.67g, 80% yield) as a solid, m.p. 86.1⁰-88.3⁰C (Lit m.p. 87-88⁰C)¹⁰




7.2 Synthesis of 3-benzo-1,3-dioxol-5-ylpropionic acid piperidide (RV-C03)

The method was adapted from that reported for piperlonguminine (section 2.2) but utilising 3-benzo-1,3-dioxol-5-ylpropionic acid and piperidine as the acid and amine components respectively. A mixture of 3-benzo-1,3-dioxol-5-ylpropionic acid (200mg, 0.0026mole, 1eq) and triethylamine (0.27ml, 0.002mole, 2eq) in dichloromethane (50ml) was stirred for 15min at 0⁰C. To this mixture methanesulfonyl chloride (0.11ml, 0.0015mole, 1.5eq) was added and stirred for further 30 min at 0⁰C. Piperidine (0.15ml, 0.0015mole, 1.5eq) was added to the mixture and stirred for 1h at 0⁰C and 1h at room temperature. Dichloromethane (50ml) was added to the mixture which was then washed with 5% HCl (3x100ml), saturated aqueous NaHCO₃ (3x100ml) and water (3x100ml). The organic fraction was dried over anhydrous sodium sulphate, filtered and rotary evaporated to yield brown oil (65% yield).

20

3-benzo-1,3-dioxol-5-ylpropionic acid piperidide (RV-C03)



¹H-NMR (CDCl₃) δ: 2.87 (t, 2H, J=7.3 CH₂), 2.57 (t, 2H, J=7.0 CH₂.CH₂) 6.70 (d, 1H J=1.5, Ar-7H), 6.72 (d, 1H J=8.0, Ar-10H), 6.66 (d, d, 1H J=1.2, 8.0 Ar-11H), 5.90 (s, 2H, O-CH₂-O), 3.55 (t, 2H, N- CH₂ (piperidine)) 3.34 (t, 2H, N- CH₂ (piperidine)) 1.62 (m, 2H, CH₂.CH₂.CH₂(piperidine)) 1.49 (m, 2H, CH₂.CH₂.CH₂(piperidine))

5 ¹³C-NMR (CDCl₃) : 25.7 (CH₂), 25.9 (CH₂), 26.6 (CH₂), 31.7 (CH₂), 35.8 (CH₂), 43.1(CH₂), 47.1(CH₂), 101.2 (CH₂), 109.2 (CH), 109.3 (CH), 121.5 (CH), 135.6 (C), 146.2 (C) 148.0 (C), 170.8 (C)

References

10 ⁹Biswanath Das., A. Kasinatham., and P. Madhusudhan. (1998). Regioselective reduction of αβ-double bond of some naturally occurring dienamides using NABH₄/I₂ system. Tetrahedron Letters **39**, 677-678.

¹⁰Perkin, Robinson, (1907) Journal of Chemical Society **91**, 1084

15 8 Synthesis of amide derivatives of methoxy-substituted cinnamic acid


A mixture of monomethoxycinnamic acid (200mg, 0.89mmol, 1eq) and triethylamine (2.4ml, 1.78mmol, 2eq) in dichloromethane (50ml) was stirred for 15min at 0⁰C. To this mixture methanesulfonyl chloride (1.02ml, 1.33mmol, 1.5eq) was added and stirred for further 30 min at 0⁰C. Piperidine (0.23ml, 1.33mmol, 1.5eq) was added to the mixture and stirred for 1h at 0⁰C and 1h at room temperature. Then dichloromethane (50ml) was added to the mixture, which was then washed with 5% HCl (3x100ml), saturated aqueous NaHCO₃ (3x100ml) and water (3x100ml). The organic fraction was dried over anhydrous sodium sulphate, filtered and rotary evaporated to yield an oil. This oil was purified by chromatography on silica gel using ethyl acetate/petroleum spirit (2:8) as an eluant.

20

25


The piperidine amide of 3,4 dimethoxycinnamic acid was prepared in the same way utilising 200mg of the acid.

1-(2-methoxy-cinnamoyl)-piperidine (RV-G01)



- ¹H-NMR (CDCl₃) δ: 7.56 (d, 1H, CH=CH), 7.29 (d, 1H, J=7.8 ArH), 7.12(d, 1H, J=7.6 ArH) 7.0 (d,d 1H, J=1.8 ArH) 6.86-6.90 (m,ArH), 6.88 (d, 1H, J=15.4 CH=CH), 3.58-3.66 (br, 4H, CH₂-N- CH₂ (piperidine)) 1.56-1.71 (m, 6H, CH₂-CH₂ CH₂ (piperidine)) 3.83 (s, 3H, OCH₃)
- ¹³C-NMR (CDCl₃) : 25.7 (CH₂), 26.0 (CH₂), 27.1 (CH₂), 43.7(CH₂), 47.4(CH₂), 55.7(CH₃), 113.4 (CH),115.3 (CH), 118.5 (CH),120.6 (CH),130.1 (CH),142.4 (CH), 137.3 (C), 160.2 (C), 165.6 (C)
- MS m/z (%) : 245(M⁺ 28), 162 (22), 161 (100), 133 (20), 118 (24), 113 (14), 84 (51)
yield 25.5%


1-(3-methoxy-cinnamoyl)piperidine (RV-G02)



- ¹H-NMR (CDCl₃) δ: 7.60 (d, 1H, J=15.4, CH=CH), 7.29 (d, 1H, J=7.8 ArH), 7.12(d, 1H, J=7.6 ArH) 7.0 (d,d 1H, J=1.8 ArH) 6.86-6.90 (m,ArH), 6.88 (d, 1H, J=15.4 CH=CH), 3.58-3.66 (br, 4H, CH₂-N- CH₂ (piperidine)) 1.56-1.71 (m, 6H, CH₂-CH₂ CH₂ (piperidine)) 3.83 (s, 3H, OCH₃)
- ¹³C-NMR (CDCl₃) : 25.7 (CH₂), 26.0 (CH₂), 27.1 (CH₂), 43.7(CH₂), 47.4(CH₂), 55.7(CH₃), 113.4 (CH),115.3 (CH), 118.5 (CH),120.6 (CH),130.1 (CH),142.4 (CH), 137.3 (C), 160.2 (C), 165.6 (C)
- MS m/z (%) : 245(M⁺ 77), 162 (65), 161 (100), 133 (20), 118 (24), 113 (14), 84 (51)


m.p. 68⁰-70⁰C, yield 31.4%

1-(4-methoxy-cinnamoyl)piperidine (RV-G03)



- 5 ¹H-NMR (CDCl₃) δ: 7.61 (d, 1H, J=15.4, CH=CH), 7.47 (d, 2H, J=7.8 ArH), 6.87-6.90 (m, 2H, ArH), 6.77 (d, 1H, J=15.4 CH=CH), 3.58-3.65 (br, 4H, CH₂-N- CH₂ (piperidine)) 1.52-1.69 (m, 6H, CH₂-CH₂ CH₂ (piperidine)) 3.82 (s, 3H, OCH₃)
¹³C-NMR (CDCl₃) : 25.6 (CH₂), 26.0 (CH₂), 26.4 (CH₂), 43.7(CH₂), 47.4(CH₂), 55.7(CH₃), 114.5 (CH), 115.6 (CH), 118.5 (CH), 121.9 (CH), 129.6 (CH), 142.2 (CH),
10 132.8 (C), 161.0 (C), 166.0 (C)
MS m/z (%) : 245(M⁺ 71), 162 (17), 161 (100), 133 (26), 118 (12), 113 (14), 84 (24), 77 (36)


1-(3,4-dimethoxycinnamoyl)piperidine (RV-G04)



- 15 ¹H-NMR (CDCl₃) 60MHz δ: 7.61 (1H, CH=CH), 7.23 (1H, ArH), 6.98(1H, ArH) 6.82 (1H, J=1.8 ArH) 6.68 (1H, CH=CH), 3.58-3.65 (br, 4H, CH₂-N- CH₂ (piperidine)) 1.5-1.8 (6H, CH₂-CH₂ CH₂ (piperidine)) 3.91 (s, 6H, OCH₃)₂
MS m/z (%) : 275(M⁺ 62), 192 (48), 191 (100), 161 (18), 118 (11), 84 (26), 77 (12), yield
20 42.3%

CLAIMS

1. A method of treating a subject having a skin condition selected from the group of conditions (a) those treatable by stimulation of melanocyte proliferation and (b) melanomas, which comprises administering to the subject an effective amount of a compound of formula (1)



(1)

- wherein $n = 0, 1$ or 2 ;
- 10 when $n = 0$, R^2 and R^3 represent hydrogen atoms or together represent a carbon to carbon double bond;
- when $n = 1$, or 2 R^2 and R^3 together and R^4 and R^5 together represent carbon to carbon double bonds, R^2 and R^3 together represent a carbon to carbon double bond and R^4 and R^5 represent hydrogen atoms, or R^2, R^3, R^4 and R^5 all represent hydrogen atoms;
- 15 $m = 1, 2$ or 3 ;
- when $m = 1$, R^1 represents an alkoxy group having from 1 to 3 carbon atoms or a hydroxy group;
- when $m = 2$, each R^1 independently represents an alkoxy group having from 1 to 3 carbon atoms or the two R^1 's together represent a 3', 4'-methylenedioxy group;
- 20 when $m = 3$, two R^1 's together represent a 3'-4'-methylenedioxy group and the other R^1 represents an alkoxy group having from 1 to 3 carbon atoms or a hydroxy group;
- R^6 represents a pyrrolidino, piperidino, 4-methylpiperidino or morpholino group, a N-monoalkylamino group of 4 to 6 carbon atoms, a N-monocycloalkylamino group of 4 to 7 carbon atoms, a 3'-4'-methylenedioxy-substituted benzylamino or 2-phenethylamino
- 25 group or R^6 represents an alkoxy group of 1 to 6 carbon atoms;

in any of its E, Z geometrically isomeric forms, but excluding compounds of formula (1) in which :

5 (a) n is 0, 1, or 2, m is 2, the R's together represent a 3', 4'-methylenedioxy group, R² and R³, together with the carbon atoms to which they are attached form a carbon to carbon double bond and, when n is 1 or 2, R⁴ and R⁵, together with the carbon atoms to which they are attached, form a carbon to carbon double bond and R⁶ is piperidino, or

10 (b) n is 1 and (i) m is 3, the R's being 3', 4'-methylenedioxy and 6'-methoxy or (ii) m is 2, the R's being 3'-hydroxy-4'-methoxy; or (iii) m is 1 and the R¹ is 4'-hydroxy; and R² to R⁶ are as defined in case (a) above, or

(c) n is 1, R⁶ is piperidino, pyrrolidino, isobutylamino or methoxy and all other symbols are defined as in case (a) above, or

15 (d) n is 1, R⁴ and R⁵ represent hydrogen atoms and either R² and R³ also represent hydrogen atoms or R² and R³ together with the carbon atoms to which they are attached form a carbon to carbon double bond; and m, R¹ and R⁶ are as defined in case (a) above;

and in all of which cases (a) to (d) the molecule is in the E,E or all E geometric configuration or in case (a) when n is 1 may be in the Z,Z, Z,E or E,Z geometric configuration.

20 2. The method of Claim 1, wherein the subject is a patient suffering from a melanoma.

25 3. The method of Claim 1, wherein the subject is a patient suffering from a skin disorder treatable by stimulation of melanocyte proliferation.

4. The method of Claim 1, wherein the skin disorder is vitiligo.

30 5. The method of Claim 1, wherein the compound is administered topically to the area of the skin to be treated.

6. The method of any preceding claim, wherein the compound of formula (1) is one in which:

5 m = 2, the R¹ groups together represent 3',4'-methylenedioxy and R⁶ represents an alkylamino group having 5 or 6 carbon atoms or a cycloalkylamino group having from 4 to 7 carbon atoms, especially a cyclohexylamino group, or R⁶ represents a 3,4-methylenedioxy-substituted benzylamino or 2-phenethylamino group.

7. The method of Claim 6, wherein the compound is of the E, E geometric configuration.

10

8. Compounds having the definition given within Claim 1, for use in treating a skin condition defined within Claim 1 which is a disorder.

15

9. Compounds according to Claim 8, wherein the skin condition is defined in Claim 2, 3 or 4.

10. Compounds according to Claim 8 or 9, for topical application to the area of the skin to be treated.

20

11. Compounds according to Claim 8, as further defined by Claim 6 or 7.

12. Each of the compounds specifically hereinbefore listed in Example 10, other than those excluded by the exclusion clause of Claim 1, for use in treating a skin condition defined within Claim 1 which is a disorder.

25



INVESTOR IN PEOPLE

Application No: GB 0101146.9
Claims searched: 1-12

Examiner: Annabel Ovens
Date of search: 26 June 2001

Patents Act 1977 Search Report under Section 17

Databases searched:

UK Patent Office collections, including GB, EP, WO & US patent specifications, in:

UK Cl (Ed.S): A5B

Int Cl (Ed.7): A61K 31/36, 31/4025, 31/45, 31/525, 31/5377

Other: Online: PAJ, EPODOC, WPI, CAS-ONLINE

Documents considered to be relevant:

Category	Identity of document and relevant passage	Relevant to claims
X	WO 00/02544 A2 (BTG INTERNATIONAL) see page 4 line 14-page 5 line 8	1-12
A	WO 96/25159 A1 (GOVERNMENT OF THE USA)	
X	WPI Abstract Accession No. 1997-359094/33 & JP 090151130 A (see abstract)	1-5, 8-10 and 12
X	WPI Abstract Accession No. 1993-357134/45 & JP 050262646 A (see abstract)	8-12

X Document indicating lack of novelty or inventive step
 Y Document indicating lack of inventive step if combined with one or more other documents of same category.
 & Member of the same patent family

A Document indicating technological background and/or state of the art.
 P Document published on or after the declared priority date but before the filing date of this invention.
 E Patent document published on or after, but with priority date earlier than, the filing date of this application.