



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>C12N 15/82, 15/29, 5/10</b> <b>A01H 5/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 91/13991</b> <b>(43) International Publication Date:</b> 19 September 1991 (19.09.91)
<b>(21) International Application Number:</b> PCT/GB91/00397 <b>(22) International Filing Date:</b> 14 March 1991 (14.03.91) <b>(30) Priority data:</b> 9005772.0                      14 March 1990 (14.03.90)                      GB <b>(71) Applicants (for all designated States except US):</b> ADVANCED TECHNOLOGIES (CAMBRIDGE) LTD. [GB/GB]; Westminster House, 7 Millbank, London SW1P 3JE (GB). AGRICULTURAL GENETICS COMPANY LTD. [GB/GB]; Unit 154/155 Cambridge Science Park, Milton Road, Cambridge CB4 4BH (GB). BIOTAL LTD. [GB/GB]; 5 Chiltern Close, Cardiff CF4 5DL (GB). BP NUTRITION LTD. [GB/GB]; Britannic House, Moor Lane, London WC2Y 9BU (GB). CIBA-GEIGY PLC [GB/GB]; Hulley Road, Macclesfield, Cheshire SK10 2NX (GB). IMPERIAL CHEMICAL INDUSTRIES PLC [GB/GB]; Imperial Chemical House, Millbank, London SW1P 3JF (GB). RHONE-POULENC LTD. [GB/GB]; Rainham Road South, Dagenham, Essex RM10 7XS (GB). SCHERING AG-ROCHEMICALS LTD. [GB/GB]; Hauxton, Cambridge CB2 5HU (GB). SHELL RESEARCH LTD [GB/GB]; 3 Savoy Place, London WC2R 0DX (GB). TWYFORD SEEDS LTD. [GB/GB]; 16 Stratford Place, London W1N 9AF (GB). UNILEVER U.K. CENTRAL RESOURCES LTD. [GB/GB]; Unilever House, Blackfriars, London EC4 (GB).		<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> LLOYD, Julie, Carol [GB/GB]; Department of Biology, University of Essex, Wivenhoe Park, Colchester CO4 3SQ (GB). JOHN, Ulrik [AU/AU]; 89 4th Avenue, Joslin, S.A. 5070 (AU). <b>(74) Agents:</b> WOODS, Geoffrey, Corlett et al.; J.A. Kemp & Co., 14 South Square, Gray's Inn, London WC1R 5LX (GB). <b>(81) Designated States:</b> AT (European patent), BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent), US.  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> PLANT PROMOTER  <b>(57) Abstract</b> <p>A plant promoter is provided, which was identified as a promoter for two proline-rich protein (PRP) genes. The promoter can be used to control the expression of genes in plant tissues.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

- 1 -

PLANT PROMOTER

The present invention relates to a plant promoter.

The major structural components of the plant cell wall in terms of quantity are polysaccharides, including  
5 cellulose, hemicellulose and pectin polymers which in certain specialised cell types, e.g. vascular/epidermal cells, are complemented by further components whose presence is adapted to the functional role of the cell. The plant cell wall of dicotyledonous plants has also been shown to  
10 contain two types of structural protein, the hydroxyproline-rich glycoproteins (HGRPs) and more recently a glycine-rich protein (GRPs) class. The HGRPs termed extensions have a characteristic S-P<sub>4</sub> repeat structure, the proline residues being hydroxylated and subsequently  
15 glycosylated.

A wide range of proline-rich proteins have now been isolated from dicotyledonous plants. Many of these have been shown to have cell wall associations although the functions remain as yet undefined. It is likely that in  
20 different cell types and in different species the PRPs present will vary.

The isolation of genomic and cDNA clones encoding PRPs has shown that this a class of proteins containing some structurally more diverse sequences. It has been found that  
25 some of the genes encoding the PRPs display a highly regulated pattern of expression at the messenger RNA level and are induced by wounding, fungal infection, auxin or during nodulation. More recently two reports have come out showing that the expression of specific PRP genes is  
30 correlated with specific developmental events.

In contrast to the position for dicot plants, very little data has been accumulated in relation to PRPs from monocot species. Purification of an extensin-like protein from maize was reported and only low levels of  
35 hydroxyproline-residues were found. Subsequent to this a highly repetitive PRP cDNA sequence from maize has been

- 2 -

published and mRNA expression studies have shown that highest levels are present in root tip and coleoptile. This protein shares a significant level of similarity with PRPs from dicot species with the exception that it contains a high proportion of threonine residues.

We have now identified a promoter sequence for each of two proline-rich protein (PRP) genes. Accordingly, the present invention provides a promoter having the nucleotide sequence:

10 (a) from -1369 to -49 upstream of the PRP140 gene,  
or

(b) from -2316 to -12 upstream of the PRP378 gene;  
the sequence optionally being modified by one or more base substitutions, insertions and/or deletions and/or by an  
15 extension at either or each end provided that the  
thus-modified sequence is capable of acting as a promoter.

The invention also provides a DNA fragment comprising such a promoter operably linked to a heterologous gene encoding a protein. Additionally provided  
20 is a vector which comprises a heterologous gene encoding a protein under the control of a promoter as above such that the gene is capable of being expressed in a plant cell transformed with the vector. A suitable vector is one in which the promoter is fused directly to the 5'-end of the  
25 gene. The vector may further contain a region which enables the gene and the promoter to be transferred to and stably integrated in a plant cell genome. The vector is generally a plasmid.

Plant cells can be transformed with such a vector.  
30 The invention therefore further provides plant cells which harbour a promoter as above operably linked to a heterologous gene encoding a protein. Transgenic plants may be regenerated from such plant cells. A transgenic plant can be obtained which harbours in its cells a promoter as  
35 above operably linked to a heterologous gene encoding a protein. Seed may be obtained from the transgenic plants.

- 3 -

The invention further provides a method of producing a desired protein in a plant cell, which method comprises:

(i) transforming a plant cell with a vector according to the invention, the protein encoded by the gene under the control of the said promoter being the desired protein; and

(ii) culturing the transformed plant cell under conditions which allow expression of the protein.

10 The invention additionally provides a method of producing a transgenic plant capable of producing a desired protein, which method comprises:

(i) transforming a plant cell with a vector according to the invention, the protein encoded by the gene under the control of the said promoter being the desired protein; and

(ii) regenerating plants from the transformed cells.

The desired proteins can be isolated from the transformed plant cells obtained by the first method and from the plants obtained by the second method.

The present promoter is composed of the sequence upstream of the wheat PRP140 gene from base -1369 to base -49 or of the wheat PRP378 gene from base -2316 to base -12, 25 base 1 being A of the ATG translational start codon for PRP. The sequence of the PRP140 promoter can be deduced from Figure 6. The sequence of the PRP378 promoter can be deduced from Figure 5.

The promoter may be obtained by preparing a genomic 30 library of wheat DNA, screening the library for the PRP140 or PRP378 gene and digesting the sequence upstream of the wheat PRP140 or PRP378 gene with appropriate restriction enzymes and/or exonucleases. There is a NspBII restriction site at base -49 and a XbaI restriction site at base -1369 35 of the upstream sequence of the PRP140 gene. There is a NotI restriction site at base +72 and a HindIII site at base

- 4 -

-2316 of the sequence of the PRP378 gene.

Several plasmid vectors have been prepared which contain an upstream sequence of the wheat PRP140 or PRP378 gene comprising the promoter sequence. These vectors 5 include pIPKH-12/2.3, pXNot/1.45, pBIXN/1.3 and pHN/2.4. E. coli MC 1022 harbouring these vectors were deposited at the National Collection of Industrial and Marine Bacteria, Aberdeen, GB on 9 November 1989 under accession numbers NCIMB 40223, NCIMB 40224, NCIMB 40225 and NCIMB 40226 10 respectively. pBIXN/1.3 was deposited under the designation pIPXN/1.3.

The PRP140 promoter may be released from pXNot/1.45 by digesting the plasmid with XbaI and NspBII. The PRP378 promoter may be released from pHN/2.4 by linearisation with 15 NotI-BstXI, incubation with a nuclease, religation of the plasmid population, selecting a plasmid with a deletion at its 3'-end to base-12 of the PRP 378 gene and digesting the plasmid with HindIII.

The promoter sequence may be modified by one or 20 more base substitutions, insertions and/or deletions and/or by an extension at either or both ends. However, the modified promoter sequence must still be capable of acting as a promoter. Sequences from base -769 to base -49 of the upstream sequence of the wheat PRP140 gene, and shorter, 25 have been found not to be sufficient to direct expression of a protein at satisfactory levels. On the other hand, the sequence from base -961 to base -49 upstream of the wheat PRP140 gene does direct protein expression at a satisfactory level. Typically there is a degree of homology of at least 30 60% between a modified sequence and the unmodified natural sequence from base -1369 to base -49 upstream of the wheat PRP140 gene or from base -2316 to -12 upstream of the wheat PRP378 gene. The degree of homology may be at least 75%, at least 85% or at least 95%.

35 A longer sequence may be provided which extends upstream of base -1369 of the PRP140 gene or of base -2316

- 5 -

of the PRP378 gene, for example to another restriction site. An extension upstream typically comprises the natural nucleotide sequence upstream of base -1369 of the PRP140 gene or of base -2316 of the PRP378 gene. Such a longer  
5 sequence may be obtained from a genomic library of wheat DNA as above.

The upstream sequence of the wheat PRP 140 gene from base -961 to base -49 may also be modified by one or more base substitutions, insertions and/or deletions and/or  
10 by an extension at either or both ends. Again, such a modified sequence must be capable of acting as a promoter. There may be a degree of homology of at least 60%, for example at least 75%, at least 85% or at least 95%, between the unmodified and modified longer sequences.

15 A modified promoter sequence may be obtained by introducing changes into the natural promoter sequence. This may be achieved by any appropriate technique, including restriction of the natural sequence with an endonuclease, insertion of linkers, use of an exonuclease and/or a  
20 polymerase and site-directed mutagenesis. A shorter DNA sequence therefore may be obtained by removing nucleotides from the 5'-terminus or the 3'-terminus of the natural promoter sequence, for example using an exonuclease such as exonuclease III or BAL 31.

25 Whether a modified sequence is capable of acting as a promoter may be readily ascertained. The modified sequence is placed upstream of a protein coding sequence, such as the bacterial reporter gene  $\beta$ -glucuronidase as in the Example. Tobacco leaf discs can then be transformed.  
30 The protein expressed by the transformed cells are then assayed, in the case of  $\beta$ -glucuronidase as described in Example 1.

The promoter may be operably linked to a heterologous gene encoding a protein. The heterologous gene  
35 may encode any protein it is desired to express. By "heterologous" is that the gene is not naturally operably

- 6 -

linked to the promoter. The gene does not therefore encode PRP140 in the case where the promoter is derived from the upstream sequence of the PRP140 gene or PRP378 in the case where the promoter is derived from the upstream sequence of the PRP378 gene. The protein may additionally comprise a transit peptide sequence at its N-terminus, encoded within the heterologous gene sequence.

The promoter is typically used to control the expression of genes in plant tissues. The protein whose expression is controlled by the promoter may be a protein encoded by a herbicide-resistance gene or a protein conferring biological control of pests or pathogens. The protein may therefore be an insecticidal protein, such as B. thuringiensis toxin, to give resistance to leaf-eating insects. Other uses to which the promoter may be put are the production of viral coat proteins to protect against viral infection, the production of high value proteins such as pharmaceuticals and the production of proteins to alter taste or nutritive value of forage grasses, etc.

The promoter sequence may be fused directly to a heterologous gene or via a linker. The linker sequence may comprise an intron. Excluding the length of any intron sequence, the linker may be composed of up to 45 bases, for example up to 30 or up to 15 bases. The linker sequence may comprise a sequence encoding a transit amino acid sequence, for example a transit sequence capable of directing a protein to a chosen subcellular locality such as the chloroplasts or mitochondria. The linker sequence may comprise a sequence having enhancer characteristics, to boost expression levels.

DNA fragments and vectors can be prepared in which the promoter is operably linked to a heterologous gene. The fragments and vectors may be single or double stranded. Plant cells can be transformed by way of such fragment directly or by way of such a vector. The vector incorporates the heterologous gene under the control of the

- 7 -

promoter. The vector contains regulatory elements capable of enabling the gene to be expressed in a plant cell transformed with the vector. Such regulatory elements include, besides the promoter, translational initiation and/or termination sequences. The vector typically contains too a region which enables the chimaeric gene and associated regulatory control elements to be transferred to and stably integrated in the plant cell genome.

The vector is therefore typically provided with transcriptional regulatory sequences and/or, if not present at the 3'-end of the coding sequence of the gene, a stop codon. A DNA fragment may therefore also incorporate a terminator sequence and other sequences which are capable of enabling the gene to be expressed in plant cells. An enhancer or other element able to increase or decrease levels of expression obtained in particular parts of a plant or under certain conditions may be provided in the DNA fragment and/or vector. The vector is also typically provided with an antibiotic resistance gene which confers resistance on transformed plant cells, allowing transformed cells, tissues and plants to be selected by growth on appropriate media containing the antibiotic.

Transformed cells are selected by growth in an appropriate medium. Plant tissue can therefore be obtained comprising a plant cell which harbours the heterologous gene under the control of the promoter, for example in the plant cell genome. The gene is therefore expressible in the plant cell. Plants can then be regenerated which include the heterologous gene and the promoter in their cells, for example integrated in the plant cell genome, such that the gene can be expressed. The regenerated plants can be reproduced and, for example, seed obtained.

A preferred way of transforming a plant cell is to use Agrobacterium tumefaciens containing a vector comprising the promoter operably linked to the heterologous gene. A hybrid plasmid vector may therefore be employed which

- 8 -

comprises:

(a) the heterologous gene under the control of the promoter and other regulatory elements capable of enabling the gene to be expressed when integrated in the genome of a plant cell;

(b) at least one DNA sequence which delineates the DNA to be integrated into the plant genome; and

(c) a DNA sequence which enables this DNA to be transferred to the plant genome.

10 Typically the DNA to be integrated into the plant cell genome is delineated by the T-DNA border sequences of a Ti-plasmid. If only one border sequence is present, it is preferably the right border sequence. The DNA sequence which enables the DNA to be transferred to the plant cell genome is generally the virulence (vir) region of a  
15 Ti-plasmid.

The heterologous gene and its transcriptional and translational control elements, including the promoter, can therefore be provided between the T-DNA borders of a  
20 Ti-plasmid. The plasmid may be a disarmed Ti-plasmid from which the genes for tumorigenicity have been deleted. The gene and its transcriptional and control elements, including the promoter, can, however, be provided between T-DNA borders in a binary vector in trans with a Ti-plasmid with a  
25 vir region. Such a binary vector therefore comprises:

(a) the heterologous gene under the control of the promoter and other regulatory elements capable of enabling the gene to be expressed when integrated in the genome of a plant cell; and

30 (b) at least one DNA sequence which delineates the DNA to be integrated into the plant genome.

Agrobacterium tumefaciens, therefore, containing a hybrid plasmid vector or a binary vector in trans with a Ti-plasmid possessing a vir region can be used to transform  
35 plant cells. Tissue explants such as stems or leaf discs may be inoculated with the bacterium. Alternatively, the

- 9 -

bacterium may be co-cultured with regenerating plant protoplasts. Plant protoplasts may also be transformed by direct introduction of DNA fragments which encode the heterologous gene and in which the promoter and appropriate  
5 other transcriptional and translational control elements are present or of a vector incorporating such a fragment.

Direct introduction may be achieved using electroporation or polyethylene glycol, microinjection or particle bombardment.

Plant cells from monocotyledonous or dicotyledonous  
10 plants can be transformed according to the present invention. Monocotyledonous species include barley, wheat, maize and rice. Dicotyledonous species include tobacco, tomato, sunflower, petunia, cotton, sugarbeet, potato, lettuce, melon, soybean, canola (rapeseed) and poplars.

15 Tissue cultures of transformed plant cells are propagated to regenerate differentiated transformed whole plants. The transformed plant cells may be cultured on a suitable medium, preferably a selectable growth medium. Plants may be regenerated from the resulting callus. Transgenic plants  
20 are thereby obtained whose cells harbour the promoter operably linked to the heterologous gene, for example integrated in their genome. The gene is consequently expressible in the cells. Seed from the regenerated plants can be collected for future use.

25 The following Examples illustrate the present invention. In the accompanying drawings:

Figure 1 shows the restriction map of the insert DNA of phage 3CX2;

Figure 2 shows the restriction map of the insert  
30 DNA of phage E3.2;

Figure 3 shows the sequence of the PRP 378 gene;

Figure 4 shows the sequence of the PRP 140 gene;

Figure 5 shows the regulatory sequence upstream of the PRP 378 gene;

35 Figure 6 shows the regulatory sequence upstream of the PRP 140 gene;

- 10 -

Figure 7 shows the construction of pBIHN/2.4;

Figure 8 shows the construction of pBIXN/1.3;

Figure 9 shows the construction of pBID -234,  
-400, -540, -769 and -961 N;

5 Figure 10 shows the construction of pIPKH -6, -12  
and -50/2.3;

Figure 11 shows the construction of pIPH-6/2.3;

Figure 12 shows the construction of pIP S-6/762 and  
pIP P-6/615;

10 Figure 13 shows the construction of pIPD-6/1839,  
1510, 1289, 1023 and 816;

Figure 14 shows the sequences of the fusion  
junctions of constructs based on BIN 19;

Figure 15 shows the sequences of the fusion  
15 junctions of constructs based on pUC19;

Figure 16 shows the results of Gus assays for the  
PRP140 promoter deletions;

Figure 17 shows the results of Gus assays for the  
PRP 378 deletions; and

20 Figure 18 shows the nucleotide sequence of the cDNA  
clone (WPRP1).

#### EXAMPLE 1

##### 1. Isolation of PRP Genes

The strategy employed to isolate wheat PRP  
25 promoters involved first constructing a genomic library of  
wheat DNA and then screening of this library using a cDNA  
probe for a PRP gene.

High molecular weight DNA was isolated from dark  
grown shoots of Triticum aestivum cv. Chinese Spring  
30 (Lazarus et al, Plant Mol. Biol. 5, 8-24, 1985). Conditions  
for partial digestion with Sau3A were established. DNA  
fragments of 18-25 kb were purified by size fractionation on  
sucrose density gradients. Lambda Charon 35 (Loenen and  
Blattner, Gene 26, 171-179, 1983) vector DNA was prepared by  
35 digestion with BamHI and purification on sucrose density  
gradients. Vector DNA (1.5  $\mu$ g) and wheat DNA (3.5  $\mu$ g) were

- 11 -

ligated at high concentration (500 ng/ $\mu$ l) using T4 DNA ligase and subsequently packaged in vitro using commercially available extracts (Stratagene).

About  $2.6 \times 10^6$  recombinants were plated onto the 5 host E. coli K803 and DNA lifts from the plaques taken onto nitrocellulose filters. The wheat PRP cDNA WPRP1 (Reference Example) was random primer labelled using  $^{32}\text{P}$  dATP and used to probe the library lifts. Positively hybridizing plaques were purified to homogeneity by several rounds of screening 10 and then DNA from these positive phage was purified by further analysis.

## 2. Characterization of the PRP Genes

Restriction maps were made of the insert DNA of the positive phages 3cX2 (Fig. 1) and E3.2 (Fig. 2) and the 15 position and orientation of the PRP genes determined by hybridization studies. DNA fragments covering these regions were subcloned into the plasmid vector pUBS1, which is a pUC19 derivative containing the polylinker region of the Bluescript plasmid of Stratagene (Raines et al, Nucl. Acids. 20 Res. 16, 7931-7942, 1988). Overlapping sequence was obtained from these clones by Exonuclease III digestion and double-stranded dideoxy nucleotide sequence analysis. Data were assembled and analysed using the Staden packages. The extent of the sequence determined is indicated on Figs 1 and 25 2.

DNA sequence comparisons between the PRP 378 gene (Fig. 3) and the cDNA probe used to isolate it revealed a few base differences which suggested that the gene copy isolated may represent an allele of the cloned mRNA. The 30 PRP 140 gene (Fig. 4) is evidently a novel form of PRP.

## 3. PRP Upstream Sequences

The regulatory sequence upstream of the PRP 378 and PRP 140 genes, including all of those used in the constructs described below, are given in Figs 5 and 6 respectively. 35 Restriction enzyme sites used during subsequent cloning procedures are indicated and underlined. It was proposed

- 12 -

that these sequences would contain all of the promoter and other regulatory elements necessary to direct correct expression of protein coding sequences placed downstream under their control.

5           In order to test this proposal and assess the usefulness of the PRP 378 and PRP 140 promoters in the context of plant transformation a series of constructs were prepared. In each case part of the PRP 378 or PRP 140 upstream regions was placed in front of the bacterial  
10 reporter gene  $\beta$ -glucuronidase and the nopaline synthase terminator sequences in a suitable vector. For the PRP 378 promoter this was a high copy number pUC19 based plasmid for transient expression in Zea mays protoplasts. For the PRP 140 promoter this was a BIN19 based vector suitable for  
15 transformation into Nicotiana tabacum. In both cases this allows expression, controlled by the PRP promoters, to be detected by an enzyme assay for  $\beta$ -glucuronidase (Gus) activity using a fluorogenic substrate in extracts from transformed plants or protoplasts.

20 (a) pIPPKGT and pBIPKGT

pIPPKGT is the EcoRI/SalI insert of pRAJ275 (Jefferson, Plant. Mol. Biol. Reporter 5, 387-405, 1987) blunted with Klenow fragment of E. coli DNA polymerase I, and ligated into the Klenow blunted Asp718 site of  
25 pUCPolyTer (the nopaline synthetase terminator fragment cloned as a SstI/EcoRI fragment into pUC19). The corresponding BIN19 based plasmid is pBIPKGT. pIPPKGT and pBIPKGT can be obtained from each other.

(b) BIN19 Based Constructs (pBI series)

30 (i) Construction of the plasmid pBIHN/2.4 containing 2318 bp of sequence 5' of the PRP 378 translation initiation codon and 24 amino acids of PRP 378 coding sequence in a translational fusion to  $\beta$ -glucuronidase (Fig. 7)

A 2.4 Kb HindIII-NotI fragment of the PRP 378 gene  
35 (Fig. 1) was subcloned into the plasmid vector pUBS1 to form the plasmid pHN/2.4. pHN/2.4 was linearised with NotI and

- 13 -

the site filled in using Klenow fragment of E. coli DNA polymerase I. The insert was excised by digestion with HindIII and inserted into HindIII-SmaI digested pBI101.3 (Jefferson, Plant. Mol. Biol. Reporter 5, 387-405, 1987) to form pBIHN/2.4. The fusion junction was checked by nucleotide sequencing and its sequence along with the predicted amino acid sequence is shown in Fig. 14.

(ii) Construction of the plasmid pBIXN/1.3, a transcriptional fusion of 1321 bp of 5' non-coding sequence from PRP 140 to  $\beta$ -glucuronidase (Fig. 8)

A 1.45 kb XbaI-NotI fragment of the PRP 140 gene (Fig. 2) was subcloned into the plasmid vector pUBS1 to form pXNot/1.45. A 1.3 kb fragment was excised by digestion with XbaI and NspBII (Fig. 8) and inserted into XbaI-SmaI digested pBIPKGT to form pBIXN/1.3. The sequence of the fusion junction is shown in Fig. 14.

(iii) Construction of the plasmids pBID -234, -400, -540, -769 and -961 N representing deletion of the PRP 140 promoter from its 5' end (Fig. 9)

The plasmid pXNot/1.45 (Fig. 8) was linearised by digestion with XbaI-PstI. Incubation with Exonuclease III for various periods resulted in deletion of varied lengths of sequence from the XbaI end. Treatment with Nuclease SI and the Klenow fragment of E. coli DNA polymerase enabled the plasmid population to be religated as closed circles. Nucleotide sequencing was employed to determine the extent of deletion from the XbaI end of a representative portion of the population.

On the basis of the information provided by sequencing, plasmids with deletion to within 234, 400, 540, 769 and 961 nucleotides 5' of the translation initiation codon of PRP 140 were chosen for evaluation of promoter activity (Fig. 6). The 5 restriction fragments were excised by digestion with HindIII and NspBII and inserted into

- 14 -

HindIII-SmaI digested pBIPKGT to form pBID -234, -400, -540, -769 and -961 N. In all 5 constructs the sequence of the fusion junction is identical to that of pBIXN/1.3 (Fig. 14).

(c) pUC19 Based Constructs (pIP series)

- 5 (i) Construction of the plasmids pIPKH -6, -12 and -50/2.3 representing deletion of the PRP 378 promoter from its 3' end (Fig. 10)

The plasmid pHN/2.4 (Fig. 7) was linearised by digestion with NotI-BstXI. Incubation with Exonuclease III  
10 for various periods resulted in deletion of varied lengths of sequence from the NotI end. Treatment with Nuclease SI and the Klenow fragment of E. coli DNA polymerase enabled the plasmid population to be religated as closed circles. Nucleotide sequencing was employed to determine the extent  
15 of deletion from the NotI end of a representative portion of the population.

On the basis of the information provided by sequencing, plasmids pHD -6, -12, -50/2.3 with inserts which extended approximately 2.3 kb downstream from the HindII  
20 sites to within 6, 12 and 50 nucleotides 5' of the translation initiation codon of PRP 378 were chosen for evaluation of promoter activity (Fig. 5). The 3 respective plasmids were linearised by a partial SstI digestion and the overhang blunted with Mung Bean Nuclease. The 3 inserts  
25 were excised by digestion with HindIII and inserted into HindIII-SmaI digested pIPPKGT to form pIPKH -6, -12, -50/2.3. The sequences of the fusion junctions are shown in Fig. 15.

(ii) Construction of pIPH-6/2.3 (Fig. 11)

30 This plasmid is similar to pIPKH-6/2.3 except that the full length PRP 378 promoter has been inserted into pBI201.1 (Jefferson, 1987) and has a different junction (Fig. 15) without a "Kozak" consensus initiating ATG.

(iii) Construction of pIP S-6/762, P-6/615 representing  
35 short forms of the PRP 378 promoter deleted from its 5' end

- 15 -

(Fig. 12)

The plasmid pIPH-6/2.3 (Fig. 11) was linearised by PstI or partial SstI digestion and the overhang blunted with Mung Bean Nuclease. Inserts with respectively 762 and 615 nucleotides of the PRP 378 promoter in a cassette with the GUS gene and the NOS terminator were excised by digestion with EcoRI and inserted into EcoRI/SmaI digested pUC19. In both constructs the sequence of the fusion junction is identical to that of pIPH-6/2.3 (Fig. 15).

10 (iv) Construction of pIPD-6/1839, 1510, 1289, 1023, 816 representing longer forms of the PRP 378 promoter deleted from its 5' end (Fig. 13)

The plasmid pHN/2.4 (Fig. 7) was linearised by digestion with HindIII-ApaI. Incubation with Exonuclease III for various periods resulted in deletion of varied lengths of sequence from the HindIII end. Treatment with Nuclease SI and the Klenow fragment of E. coli DNA polymerase enabled the plasmid population to be religated as closed circles. Nucleotide sequencing was employed to determine the extent of deletion from the HindIII end of a representative portion of the population.

On the basis of information provided by sequencing plasmids pD-1839, -1510, -1289, -1023 and -816 N with inserts extending from 1839 to 816 nucleotides 5' of the translation initiation codon of PRP 378 were digested with KpnI-SstII. The excised fragments were ligated with EcoRI-partial SstII digested pIPH-6/2.3 (providing the proximal portion of the PRP 378 promoter in conjunction with the GUS gene and the NOS terminator) which in turn was ligated into EcoRI-KpnI digested pUC19. Once again in all constructs the sequence of the fusion junction is identical to that of pIPH-6/2.3 (Fig. 15).

#### 4. Introduction of PRP Promoter Fusions into Nicotiana

Constructs were mobilised from Escherichia coli MC1022 into Agrobacterium tumefaciens LBA4404 as described (Bevan, Nucl. Acids. Res. 12, 8711-8721, 1984). Leaf discs

- 16 -

of Nicotiana tabacum var. Samsun were transformed as described (Horsch et al, Science 223, 496-498, 1984) and selected on shooting medium containing 100 µg/ml kanamycin. Typically between 5 and 15 plants containing each construct 5 were regenerated and assayed.

#### 5. β-Glucuronidase Assay of Transformed Plants

The activity of the PRP promoters in individual transformants was determined by measuring β-glucuronidase activity in leaf extracts. Tissue extracts were prepared 10 and analysed for fluorescence of the reaction product 4-methyl umbelliferone as described (Jefferson, 1987). Reactions were usually incubated at 37°C for 4 hours with aliquots sampled at 2 hour intervals. The protein concentration in each extract was measured to allow direct 15 comparisons to be made between them (using a Bio-Rad kit).

Results of assays on PRP140 promoter derived constructs expression in nmol 4-methyl umbelliferone produced/min/mg protein are shown in Fig. 16. Because of the variation commonly observed between individual 20 transformed plants it is difficult to ascertain whether dissection of the PRP 140 promoter from 1369 to 961 nucleotides in length leads to a significant loss of activity. However, it is clear that the 769 and 540 nucleotide forms have substantially less activity and that 25 the 400 and 234 nucleotide forms are effectively inactive.

#### 6. Introduction of PRP 378 Promoter Fusions into Protoplasts Derived from a Suspension Culture of Zea mays Anther Cells

The pUC19 based constructs were assayed for 30 transient expression by Ca<sup>++</sup>- PEG mediated transformation of Zea mays anther derived cell suspension (cell line Ba Tan Bai) protoplasts based on the method of Chen et al. (Chen, D.F., Tassie, A., Dale, P.J., Goldsbrough, A.P., Liang, W., Bevan, M.W., Flavell, R.B. and Xia, Z.A. (1988). In: Puite, 35 K.J., Dons, J.J.M., Huizing, H.J., Kool, A.J., Koornneef, M. and Krens, F.A. (eds) Progress in Plant Protoplast Research.

- 17 -

Kluwer Academic Publishers, Dordrecht). Typically for each construct 3 replicates of 50  $\mu\text{g}$  of DNA and  $1 \times 10^6$  protoplasts were performed. As controls protoplasts were also transformed with identical amounts of herring sperm DNA and the plasmid CG20 carrying one copy of the 850 bp form of the cauliflower mosaic virus (CaMV) 35S promoter.

7.  $\beta$ -Glucuronidase Assay of PEG mediated Transformed Zea mays Protoplasts

Lysates of transformed protoplasts were assayed for  $\beta$ -glucuronidase in the same manner as that for transformed plants (Jefferson, 1987). Expressing activity per  $1 \times 10^4$  protoplasts allowed comparisons to be made between constructs tested in the same experiment.

Results of assays on PRP 378 promoter derived constructs expressed in nmol 4-methyl umbelliferone produced/min/ $1 \times 10^4$  protoplasts are shown in Table 1 and Fig. 17.

Table 1. Gus Assays: PRP378 Promoter 3' Deletions

<u>Construct</u>	<u>nMOL of 4-MU/min/<math>1 \times 10^4</math> protoplasts</u>
20 CG20 (CaMV 35S)	$3.59 \pm 2.96$ (100%)
Herring sperm DNA	0.0308 (0.858)
pipKH -6 /2.3	$9.65 \pm 0.63$ (269)
-12	$11.92 \pm 4.22$ (332)
-50	$4.73 \pm 0.66$ (132)

25 The 2.3 kbp form of the PRP 378 promoter is significantly stronger than the 850 bp form of the CaMV35S promoter in maize protoplasts (Table 1). In addition it is sensitive to relatively short deletions from its 3' end. Thus pipKH-12/2.3, the construct that places the promoter in  
30 almost its native context (Fig. 15) with respect to the initiating "ATG" of the GUS gene, exhibits highest levels of expression; whilst activity declines if, in effect, the promoter is moved (Fig. 15) further or closer to the ATG.

Figure 17 illustrates the effect of deletions from

- 18 -

5' end of the PRP 378 promoter. A deletion of 800 bp reduces activity to less than 60% of the full length form, and activity continues to decline with extent of deletion. Although the full length form of the PRP 378 promoter in this experiment does not have the same level of activity with respect to CaMV 35S as those in Table 1, its fusion junction is different and the initiating ATG of the  $\beta$ -glucuronidase gene does not have the "Kozak" consensus (Fig. 15).

#### 10 REFERENCE EXAMPLE

A cDNA, designated WPRP1, encoding a wheat PRP has been isolated and sequenced. The WPRP1 clone was detected whilst a wheat cDNA library was being screened using antibodies raised against chloroplastic fructose-1,5-bisphosphatase. Why this antibody should cross-react with the WPRP1 clone is not known. The cDNA isolated initially was not full length (1.3 Kb). The complete sequence came from a longer cDNA obtained by rescreening the cDNA library using the shorter clone.

20 The complete nucleotide sequence of the cDNA clone (WPRP1) from wheat with its deduced amino acid sequence is shown in Figure 18. This 1548 bp cDNA contains the complete coding sequence of 1137 base pairs with 101 bp of 5' non-coding sequence and 310 bp of 3' non-coding sequence  
25 terminating in a poly-adenylation tail.

#### EXAMPLE 2

The localisation of Gus expression in tissues of the transformed tobacco plants of Example 1.4 was investigated. Assays were conducted as described in Example 1.5 except that tissues other than leaf were also assayed. The results are shown in Table 2 below. The values shown are pmol/min/ug protein. The numbers in brackets refer to ranking of tissue expression level within each individual transformed plant. There are significant

- 19 -

levels of expression in all tissues. The levels of expression in petioles and stems compare favourably with CaMV 35S.

Table 2

5		<u>leaf</u>	<u>petiole</u>	<u>root</u>	<u>stem</u>
PRP140: 961	3	7.93 (5)	102.75 (1)	18.82 (3)	16.96 (4)
	4	10.38 (3)	170.16 (1)	5.29 (4)	63.87 (2)
	6	6.28 (4)	65.02 (1)	5.65 (5)	30.53 (2)
1369	2	17.22 (3)	95.45 (1)	3.29 (5)	51.86 (2)
10	5	4.58 (3)	104.64 (1)	2.04 (5)	24.52 (2)
	8	16.60 (3)	305.35 (1)	7.32 (5)	40.87 (2)
	12	6.64 (5)	76.25 (1)	6.65 (4)	33.20 (2)
CaMV		39.96 (4)	58.25 (2)	50.14 (3)	62.43 (1)
BIN(0)		0.35	1.31	3.62	0.67
15		<u>apical</u>			
PRP140: 961	3	19.55 (2)			
	4	1.96 (5)			
	6	9.08 (3)			
1369	2	17.21 (4)			
20	5	4.31 (4)			
	8	26.70 (4)			
	12	12.41 (3)			
CaMV		32.48 (5)			
BIN(0)		0.32			

- 20 -

CLAIMS

1. A promoter having the nucleotide sequence:
  - (a) from -1369 to -49 upstream of the PRP140 gene, or
  - (b) from -2316 to -12 upstream of the PRP 378 gene;
- 5 the sequence optionally being modified by one or more base substitutions, insertions and/or deletions and/or by an extension at either or each end provided that the thus-modified sequence is capable of acting as a promoter.
2. A promoter according to claim 1, having a
- 10 sequence from -961 to -49 upstream of the PRP140 gene, optionally modified as specified in claim 1.
3. A DNA fragment comprising a promoter as claimed in claim 1 or 2 operably linked to a heterologous gene encoding a protein.
- 15 4. A vector which comprises a heterologous gene, encoding a protein, under the control of a promoter as claimed in claim 1 or 2, such that the gene is capable of being expressed in a plant cell transformed with the vector.
5. A vector according to claim 4, wherein the
- 20 promoter is fused directly to the 5'-end of the said gene.
6. A vector according to claim 4 or 5, which further contains a region which enables the gene and the promoter to be transferred to and stably integrated in a plant cell genome.
- 25 7. A vector according to any one of claims 4 to 6, which is a plasmid.
8. A plant cell which has been transformed with a vector as claimed in any one of claims 4 to 7.
9. A plant cell which harbours a promoter as
- 30 claimed in claim 1 or 2 operably linked to a heterologous gene encoding a protein.
10. A transgenic plant which has been regenerated from plant cells as claimed in claim 8 or 9.
11. A transgenic plant which harbours in its cells
- 35 a promoter as claimed in claim 1 or 2 operably linked to a heterologous gene encoding a protein.

- 21 -

12. Seed obtained from a transgenic plant as claimed in claim 10 or 11.

13. A method of producing a desired protein in a plant cell, which method comprises:

5 (i) transforming a plant cell with a vector as claimed in any one of claims 4 to 7, the protein encoded by the gene under the control of the said promoter being the desired protein; and

10 (ii) culturing the transformed plant cell under conditions which allow expression of the protein.

14. A method of producing a transgenic plant capable of producing a desired protein, which method comprises:

15 (i) transforming a plant cell with a vector as claimed in any one of claims 4 to 7, the protein encoded by the gene under the control of the said promoter being the desired protein; and

(ii) regenerating plants from the transformed cells.

Fig. 1.




Fig. 2.




Fig. 3.

2319 ATGGCGAGGCACAGCCTCCTTGGCCGTGCTCCTCGTGGGCTAGTCGGGCTCCGGCTTC 2378  
M A R H S L L A V L L V G L V A A S G F

2379 AGCCAGGCCGCCGCTGGCCCTGGCTGAGAGCTCCCGGAGCCGGAGCCCAAG 2438  
S Q A A A A G R G L A E K L P E P E P K  
NotI

2439 CCGACGCCGTACCCGGAGCCCAAGCCGGAACCAAGCCAGCCCAATGCCTAAGCCTGAA 2498  
P T P Y P E P K P E P K P E P M P K P E

2499 CCGATGCCAAAGCCAGAGCCTAAGCCTGAACCCATGCCAAAGCCAGAGCCTAAGCCTATG 2558  
P M P K P E P K P E P M P K P E P K P M

2559 CCCAAACCTGAGCCCAATGCCAAAGCCAGAGCCCAACCGGAACCGGAGCCCGATG 2618  
P K P E P M P K P E P K P E P K P E P M

2619 CCAAAGCCCGAACCAAACTGAGCCCAAGCCTTACCCGATGCCCAAACTGAGCCTAAG 2678  
P K P E P K P E P K P Y P M P K P E P K

2679 CCTGAGCCCAAGCCCGAGCCTATGCCAAACCTGAACCAAGCCAGAGCCCAAGCCCGAG 2738  
P E P K P E P M P K P E P K P E P K P E

2739 CCTATGCCAAACCTGAACCAAGCCCGAGCCCAAGCCTGAGCCTATGCCAAACCTGAG 2798  
P M P K P E P K P E P K P E P M P K P E

2799 CCTAAGCCGATGCCAAACCCGAAACCGAAGCCAGAGCCCAAGCCTGAGCCGATGCCCTAAG 2858  
P K P M P K P E P K P E P K P E P K P E P M P K

2859 CCAGAACCGAAGCCAGAGCCCAAGCCTGAGCCTATGCCAAACCTGAACCAAGCCAGAG 2918  
P E P K P E P K P E P M P K P E P K P E

2919 CCCAAACTCGAGCCGATGAAGCCCGAGCCTAAGCCGGTCCAAACCCGAACTAAGCCA 2978  
P K L E P M K P E P K P V P K P E P K P

2979 GATCCCAAGCCTGAGCCGATGCCAAACCCGAGCCCAAGCCAGAGCCCAAGCCTGAGCCG 3038  
D P K P E P M P K P E P K P E P K P E P K P E P

Fig.3(cont.)

3039 ATGCCAAAACCCGAACCGAAGCCGGAGCCCAAGCCCTTACCCAAATGCCCAACCTGAGCCT 3098  
 M P K P E P K P E P K P E P K P Y P M P K P E P  
 3099 AAGCCTGAGCCCAAGCCCTGAGCCGATGCCCAAAACCAGAACCAAGCCAGAGCCCAACCC 3158  
 K P E P K P E P M P K P E P K P E P K P E P K P  
 3159 GAGCCGATGAAGCCAGAGCCTAAGCCAAATGCCAAAGCCGGAACCGAAGCCAGAGCCCAAG 3218  
 E P M K P E P K P M P K P E P K P E P K P E P K  
 3219 CCTGAGCCGATGCCCAAGCCGAGCCCAAGCCCTATGCCGGAAGCCAGAGCCCAACCCAGAG 3278  
 P E P M P K P E P K P M P K P E P K P E P K P E  
 3279 CCGATGCCAAAGCCAGAGCCCTAAGCCCGAACCCATTGCCCTAAACCAGAGCCTAAGCCCGAA 3338  
 P M P K P E P K P E P L P K P E P K P E  
 3339 CCAATGCCCTAAACCAGAGCCCAAGCCCTGAACCTATGCCCTAAGCCGGAGCCCAAGCCCTGAA 3398  
 P M P K P E P K P E P M P K P E P K P E  
 3399 CCGGAACCAAGCCAGAGCCACCTCCGAAGGGCAAGCCGCCAATGACTGACAAATGATGT 3458  
 P E P K P E P P P K G K P P M T D N \*  
 3459 GATACTCACATATGACAGCTGAAGGAGGAGATCGACCCCGTCCGGAGCCACGGTGTGCTG 3518  
 XbaI  
 3519 TTTCTAGA 3526



*Fig. 4 (cont.)*

	550		570	590	
541	CGATGCATCTTTCTCGTATTCTATGTATTTTCGGTTACATTTATAAAGTTTCTGTTCATGCCA				600
	610		630	650	
601	TCCATGATGAGTATTACTACTTTGTAATTGTTATTTGCTTGGTGGTGAAGTTCAATTTT				660
	670		690	710	
661	GTACGTGACTCGTTGTTCTATTGTTGTTATCCGCATGAATGAGTTATTTGTCTAGA				715
					<u>· XbaI</u>



7/20


Fig.6.




XbaI

1    1    2    3    4    5    6    7    8    9    10   11   12   13   14   15   16   17   18   19   20   21   22   23   24   25   26   27   28   29   30   31   32   33   34   35   36   37   38   39   40   41   42   43   44   45   46   47   48   49   50   51   52   53   54   55   56   57   58   59   60   61   62   63   64   65   66   67   68   69   70   71   72   73   74   75   76   77   78   79   80   81   82   83   84   85   86   87   88   89   90   91   92   93   94   95   96   97   98   99   100   101   102   103   104   105   106   107   108   109   110   111   112   113   114   115   116   117   118   119   120   121   122   123   124   125   126   127   128   129   130   131   132   133   134   135   136   137   138   139   140   141   142   143   144   145   146   147   148   149   150   151   152   153   154   155   156   157   158   159   160   161   162   163   164   165   166   167   168   169   170   171   172   173   174   175   176   177   178   179   180   181   182   183   184   185   186   187   188   189   190   191   192   193   194   195   196   197   198   199   200   201   202   203   204   205   206   207   208   209   210   211   212   213   214   215   216   217   218   219   220   221   222   223   224   225   226   227   228   229   230   231   232   233   234   235   236   237   238   239   240   241   242   243   244   245   246   247   248   249   250   251   252   253   254   255   256   257   258   259   260   261   262   263   264   265   266   267   268   269   270   271   272   273   274   275   276   277   278   279   280   281   282   283   284   285   286   287   288   289   290   291   292   293   294   295   296   297   298   299   300   301   302   303   304   305   306   307   308   309   310   311   312   313   314   315   316   317   318   319   320   321   322   323   324   325   326   327   328   329   330   331   332   333   334   335   336   337   338   339   340   341   342   343   344   345   346   347   348   349   350   351   352   353   354   355   356   357   358   359   360   361   362   363   364   365   366   367   368   369   370   371   372   373   374   375   376   377   378   379   380   381   382   383   384   385   386   387   388   389   390   391   392   393   394   395   396   397   398   399   400   401   402   403   404   405   406   407   408   409   410   411   412   413   414   415   416   417   418   419   420   421   422   423   424   425   426   427   428   429   430   431   432   433   434   435   436   437   438   439   440   441   442   443   444   445   446   447   448   449   450   451   452   453   454   455   456   457   458   459   460   461   462   463   464   465   466   467   468   469   470   471   472   473   474   475   476   477   478   479   480   481   482   483   484   485   486   487   488   489   490   491   492   493   494   495   496   497   498   499   500   501   502   503   504   505   506   507   508   509   510   511   512   513   514   515   516   517   518   519   520   521   522   523   524   525   526   527   528   529   530   531   532   533   534   535   536   537   538   539   540   541   542   543   544   545   546   547   548   549   550   551   552   553   554   555   556   557   558   559   560   561   562   563   564   565   566   567   568   569   570   571   572   573   574   575   576   577   578   579   580   581   582   583   584   585   586   587   588   589   590   591   592   593   594   595   596   597   598   599   600   601   602   603   604   605   606   607   608   609   610   611   612   613   614   615   616   617   618   619   620   621   622   623   624   625   626   627   628   629   630   631   632   633   634   635   636   637   638   639   640   641   642   643   644   645   646   647   648   649   650   651   652   653   654   655   656   657   658   659   660   661   662   663   664   665   666   667   668   669   670   671   672   673   674   675   676   677   678   679   680   681   682   683   684   685   686   687   688   689   690   691   692   693   694   695   696   697   698   699   700   701   702   703   704   705   706   707   708   709   710   711   712   713   714   715   716   717   718   719   720   721   722   723   724   725   726   727   728   729   730   731   732   733   734   735   736   737   738   739   740   741   742   743   744   745   746   747   748   749   750   751   752   753   754   755   756   757   758   759   760   761   762   763   764   765   766   767   768   769   770   771   772   773   774   775   776   777   778   779   780   781   782   783   784   785   786   787   788   789   790   791   792   793   794   795   796   797   798   799   800   801   802   803   804   805   806   807   808   809   810   811   812   813   814   815   816   817   818   819   820   821   822   823   824   825   826   827   828   829   830   831   832   833   834   835   836   837   838   839   840   841   842   843   844   845   846   847   848   849   850   851   852   853   854   855   856   857   858   859   860   861   862   863   864   865   866   867   868   869   870   871   872   873   874   875   876   877   878   879   880   881   882   883   884   885   886   887   888   889   890   891   892   893   894   895   896   897   898   899   900   901   902   903   904   905   906   907   908   909   910   911   912   913   914   915   916   917   918   919   920   921   922   923   924   925   926   927   928   929   930   931   932   933   934   935   936   937   938   939   940   941   942   943   944   945   946   947   948   949   950   951   952   953   954   955   956   957   958   959   960   961   962   963   964   965   966   967   968   969   970   971   972   973   974   975   976   977   978   979   980   981   982   983   984   985   986   987   988   989   990   991   992   993   994   995   996   997   998   999   1000   1001   1002   1003   1004   1005   1006   1007   1008   1009   1010   1011   1012   1013   1014   1015   1016   1017   1018   1019   1020   1021   1022   1023   1024   1025   1026   1027   1028   1029   1030   1031   1032   1033   1034   1035   1036   1037   1038   1039   1040   1041   1042   1043   1044   1045   1046   1047   1048   1049   1050   1051   1052   1053   1054   1055   1056   1057   1058   1059   1060   1061   1062   1063   1064   1065   1066   1067   1068   1069   1070   1071   1072   1073   1074   1075   1076   1077   1078   1079   1080   1081   1082   1083   1084   1085   1086   1087   1088   1089   1090   1091   1092   1093   1094   1095   1096   1097   1098   1099   1100   1101   1102   1103   1104   1105   1106   1107   1108   1109   1110   1111   1112   1113   1114   1115   1116   1117   1118   1119   1120   1121   1122   1123   1124   1125   1126   1127   1128   1129   1130   1131   1132   1133   1134   1135   1136   1137   1138   1139   1140   1141   1142   1143   1144   1145   1146   1147   1148   1149   1150   1151   1152   1153   1154   1155   1156   1157   1158   1159   1160   1161   1162   1163   1164   1165   1166   1167   1168   1169   1170   1171   1172   1173   1174   1175   1176   1177   1178   1179   1180   1181   1182   1183   1184   1185   1186   1187   1188   1189   1190   1191   1192   1193   1194   1195   1196   1197   1198   1199   1200   1201   1202   1203   1204   1205   1206   1207   1208   1209   1210   1211   1212   1213   1214   1215   1216   1217   1218   1219   1220   1221   1222   1223   1224   1225   1226   1227   1228   1229   1230   1231   1232   1233   1234   1235   1236   1237   1238   1239   1240   1241   1242   1243   1244   1245   1246   1247   1248   1249   1250   1251   1252   1253   1254   1255   1256   1257   1258   1259   1260   1261   1262   1263   1264   1265   1266   1267   1268   1269   1270   1271   1272   1273   1274   1275   1276   1277   1278   1279   1280   1281   1282   1283   1284   1285   1286   1287   1288   1289   1290   1291   1292   1293   1294   1295   1296   1297   1298   1299   1300   1301   1302   1303   1304   1305   1306   1307   1308   1309   1310   1311   1312   1313   1314   1315   1316   1317   1318   1319   1320   1321   1322   1323   1324   1325   1326   1327   1328   1329   1330   1331   1332   1333   1334   1335   1336   1337   1338   1339   1340   1341   1342   1343   1344   1345   1346   1347   1348   1349   1350   1351   1352   1353   1354   1355   1356   1357   1358   1359   1360   1361   1362   1363   1364   1365   1366   1367   1368   1369   1370   1371   1372   1373   1374   1375   1376   1377   1378   1379   1380   1381   1382   1383   1384   1385   1386   1387   1388   1389   1390   1391   1392   1393   1394   1395   1396   1397   1398   1399   1400   1401   1402   1403   1404   1405   1406   1407   1408   1409   1410   1411   1412   1413   1414   1415   1416   1417   1418   1419   1420   1421   1422   1423   1424   1425   1426   1427   1428   1429   1430   1431   1432   1433   1434   1435   1436   1437   1438   1439   1440   1441   1442   1443   1444   1445   1446   1447   1448   1449   1450   1451   1452   1453   1454   1455   1456   1457   1458   1459   1460   1461   1462   1463   1464   1465   1466   1467   1468   1469   1470   1471   1472   1473   1474   1475   1476   1477   1478   1479   1480   1481   1482   1483   1484   1485   1486   1487   1488   1489   1490   1491   1492   1493   1494   1495   1496   1497   1498   1499   1500

8/20


Fig. 7.






-  PRP378 Promoter
-  GUS
-  NOS TER

9/20


Fig. 8.






-  PRP140 Promoter
-  GUS
-  NOS TER

10/20


Fig. 9.



-  PRP140 Promoter
-  GUS
-  NOS TER


11/20

Fig. 10.




12/20

Fig.11.



13/20

Fig. 12.






-  PRP378 Promoter
-  GUS
-  NOS TER

Fig.13.






Fig. 14.

CONSTRUCT BASED ON :-




CONSTRUCTS BASED ON :-



16/20

CONSTRUCTS BASED ON:



CONSTRUCTS BASED ON:





Fig. 15.

17/20

Fig. 16.



18/20




Fig. 17.

Fig. 18.

1 GAATTGGGGACACCCCATCCACTTTATCTCAGTTGAGAGGTTTAGAAGCGGGGTGAAGCACAAGTGCAGCAAGAGC  
80 TGGGAGAAGGAGCGAGCAATGGCGAGGCACAGCCCTCCCTGCGTGTCCCTCGTGGGGTGGTCCGGCCCTCCGGCTT  
M A R H S L L A V L L V G L V A A S G F  
81  
161 CAGCCAGGGCCCGTGGCCGGGCTTGTGAGAACTCCCGAGCCGAGCCCAAGCCGACCGCGTACCCAGAGC  
S Q A A A G R G L A E K L P E P E P K P T P Y P E P  
161  
241 CCAAGCCGXAACCCAGCAGCCAAATGCCCTAAGCCTGAACCCATGCCAAAGCCAGAGCCTAAGCCTTGCCCTAAACCT  
K P X P K P E P M P K P E P M P K P E P K P L P K P  
241  
321 GAACCCATGCCAAAACCCAGAGCCTAAGCCTTGCCAAAACCTGAACCCATGCCAAAAGCCAGAGCCCAAAACCCGGAGCCGAA  
E P M P K P E P K P L P K P E P M P K P E P K P E P K P E P K  
321  
401 GCCGGAGCCGATCCCAAAGCCCGAACCAAGCCTGAGCCCAAGCCTGACCCGATGCCCAAACTGAGCCTAAGCCTGAGC  
P E P I P K P E P K P E P K P D P M P K P E P K P E P K P E P  
401  
481 CCAAGCCAGAGCCGATGCCAAAACCCGGAACCAAGCCAGAGCCCAAGCCGAGCCTATGCCAAAACCTGAACCCAAAGCCCC  
K P E P M P K P E P K P E P K P E P M P K P E P K P E P K P  
481  
561 GAGCCAAAACCCGAGCCGATGAAGCCTGAGCCTAAGCCGATGCCAAAACCCGAAACCCGAAACCCAGAGCCCAAGCCGAGCC  
E P K P E P M K P E P K P M P K P E P K P E P K P E P K P E P  
561  
641 GATGCCCTAAACCAGAACCAAGCCAGAGCCCAACCCGAGCCGATAAAGCCTGAGCCTAAGCCGATGCCGAAACCCGAAAC  
M P K P E P K P E P K P E P I K P E P K P E P M P K P E P  
641  
721 CTAAGCCAGAGCCCAAGCCTGAGCCGATGCCAAAACCCGAAACCCGAAACCCGAAACCCGAGCCTGAGCCGATGCCAAAACCCG  
K P E P K P E P M P K P E P K P E P K P E P K P E P M P K P  
721  
801 GAGCCAAAACCCGAGCCGATGCCAAAACCCGAAACCCGAAACCCGAAACCCGAAACCCGAAACCCGAAACCCGAAACCCGAA  
E P K P E P K P E P M P K P E P K P E P K P E P K P Y P M P K  
801  
881 ACCTGAGCCCTAAGCCCTGAGCCCAACCCGAGCCGATGCCAAAACCCAGAACCCAGAACCCGAGCCCAAAACCCGAGCCGATGC  
P E P K P E P K P E P M P K P E P K P E P K P E P K P E P M P  
881


Fig. 18(cont.)

961	CAAGCCGGAAACCAAGCCAGAGCCCAAAACCCGAGCCCAAGCCAGAGCCCGGATGCCAAAGCCGGAAACCAAGCCCGAGCCCC	1040
	K P E P K P E P K P E P K P E P M P K P E P K P E P K P E P	
1041	AAGCCGAGCCCAATGCCGAGCCAGAAACCGAAGCCAGAGCCCAAGCCCTGAGCCGATGCCAAAGCCAGAGCCCTAAGCCCAA	1120
	K P E P M P K P E P K P E P K P E P M P K P E P K P E P K P K	
1121	ACCATTGCCCTAAACCCAGAGCCCTAAGCCCTAAGCCCTAAGCCAGAGCCCAAGCCCTGAGCCCGAACCCGAAAGCCGGAGC	1200
	P L P K P E P K P E P M P K P E P K P E P K P E P E P K P E P	
1201	CGCCTCCGAAAGGGCAAGCCACCGATGACTGACAAATTGATGTGATACTCACATATGCACAGCTGAAGGGAGGAGATCGACCCCC	1280
	P P K G K P P M T D N	
1281	GTCCGGAGCCACGATGTGCTATTTCTAGAAATAAGTGGCATAAGTATCCGGTTAGCCGAGATAGTGATGATGCATCTTTTGT	1360
	ATTCCTTGATTCACCAATTCCTTTTAGTTCTGTGTTCCATGCACCCATGATGAGTACTACTTCTGTAATTAATCAT	1440
1441	TTGCGTGTGGTATATGTTTCATCTGTGCACATGACTCAGTTGTTCTTTCGTTGATACACATGCCTTTTATTTCGTTTGCA	1520
1521	TGAAAGACACATTTTGTCCACCGAATTC	1548

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 91/00397

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>6</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC <sup>5</sup> : C 12 N 15/82, C 12 N 15/29, C 12 N 5/10, A 01 H 5/00		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
IPC <sup>5</sup>	C 12 N, A 01 H	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT <sup>9</sup></b>		
Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
A	Plant Molecular Biology, volume 11, 1988 Kluwer Academic Publishers, (Dordrecht NL), V. Stiefel et al.: "Molecular cloning of cDNAs encoding a putative cell wall protein from Zea mays and immunologi- cal identification of related polypep- tides", pages 483-493 see the whole document  ---	1-14
A	Journal of Cellular Biochemistry, Suppl. 13D, 1-7 April 1987, Alan R. Liss, Inc., (New York, US), M.P. Vallés et al.: "Structure and expression of two families of develop- mentally regulated genes from Zea mays", page 286 see abstract M241  ---	1-14
	./.	
<p><sup>10</sup> Special categories of cited documents: <sup>10</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p>		
<b>IV. CERTIFICATION:</b>		
Date of the Actual Completion of the International Search	Date of Making of this International Search Report	
14th June 1991	19. 07. 91	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE		
	Mme Dagmar FRANK	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, " with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	Chemical Abstract, volume 112, 1990, (Columbus, Ohio, US), Keller, Beat et al.: "Specific expres- sion of a novel cell wall hydroxy- proline-rich glycoprotein gene in lateral root initiation", see page 209, abstract no. 113287y & Genes Dev. 1989, 3(10), 1639-46  -----	1-14